

Efficacy of eDNA vs Conventional Monitoring Methods

This Guidance is publicly available for readers to review and use. Any comments that improve the content or workability of the guidance are welcomed by the JIP. We will endeavor to address comments in future updates to the guidance, which are planned every 1 to 2 years to capture the rapid advances in this field. Please send comments to info@iogp-edna.org.

Prepared by NatureMetrics and ID-Gene for the International Oil and Gas Producers' Association Joint Industry Program on Environmental Genomics.

CC BY-NC ([Deed - Attribution-NonCommercial 4.0 International - Creative Commons](https://creativecommons.org/licenses/by-nc/4.0/))



**NATURE
METRICS**
DNA-BASED MONITORING

ID-GENE
ecodiagnosics

Disclaimer

Whilst every effort has been made to ensure the accuracy of the information contained in this publication and any associated material, neither IOGP nor any of its Members past present or future warrants its accuracy or will, regardless of its or their negligence, assume liability for any foreseeable or unforeseeable use made thereof, which liability is hereby excluded. Consequently, such use is at the recipient's own risk on the basis that any use by the recipient constitutes agreement to the terms of this disclaimer. The recipient is obliged to inform any subsequent recipient of such terms.

Please note that this publication and any associated material is provided for informational purposes and adoption of any of its recommendations is at the discretion of the user. Except as explicitly stated otherwise, this publication must not be considered as a substitute for government policies or decisions or reference to the relevant legislation relating to information contained in it.

Where the publication contains a statement that it is to be used as an industry standard, IOGP and its Members past, present, and future expressly disclaim all liability in respect of all claims, losses or damages arising from the use or application of the information contained in this publication in any industrial application.

Any reference to third party names is for appropriate acknowledgement of their ownership and does not constitute a sponsorship or endorsement.

Revision history

RELEASE	NAME	COMPANY	DATE OF ISSUE	COMMENTS
R1	Ed Wort	NatureMetrics	2024-Jan-26	First release.
R2	Jan Pawlowski	ID-gene	2024-Nov-03	Revision of initial release to improve relevance, completeness, accuracy and readability.
R3	Ed Wort	NatureMetrics	2024-May-28	Revisions following JIPs comments
R4	Nicolas Tsesmetzis	Shell	2024-Oct-11	Addition of Disclaimer.
R5	Michael Marnane	Chevron	2024-Nov-21	Addition of Instructions to readers
R6	Michael Wright	eDNAtec	2025-Jun-18	Cross-referencing and alignment in terminology between chapters

Executive Summary

Environmental DNA (eDNA) based methods are increasingly used for biodiversity monitoring across a range of habitats in which the energy industry operates. This publication reviews eDNA and conventional biomonitoring methods, compares their efficiency, and discusses their advantages and limitations. It highlights that, although eDNA methods are still developing, there are many potential applications throughout the life cycle of an energy project, either as stand-alone or complementary monitoring tool.

This guidance consists of seven sections: (1) an introduction to the importance of environmental monitoring in the energy industry, (2) an overview of conventional monitoring methods, (3) and overview of eDNA-based methods, (4) a comparison of the efficacy of conventional and eDNA-based methods, using a selection of case studies, (5) the current application of eDNA-based methods in the energy industry, (6) considerations when deciding whether to use eDNA-based methods, and (7) future directions for eDNA-based technology. The guidance considers different habitats (freshwater, marine, terrestrial) and wide range of taxonomic groups, from vertebrates to microbes.

Based on a critical review of more than 200 scientific papers, the guidance shows that eDNA methods are mature enough to be considered as an integral part of biomonitoring in several energy industry applications. Their efficacy is particularly high in the domain of detection and identification of individual species. They also perform well in holistic biodiversity surveys, as they can detect small-sized and inconspicuous taxa, that are difficult to identify using conventional methods. On the other hand, the eDNA approach is currently less successful in inferring the abundance of organisms, especially larger-sized taxa. The taxonomic composition of communities inferred from eDNA and conventional methods is often different. This makes the eDNA approach more challenging when applied to morphology-based ecological indices, but solutions have been proposed in the literature to overcome these limitations.

The main advantages of an eDNA approach over conventional monitoring methods are its lower level of effort and cost, its ability to detect species when they are difficult to observe directly, and to identify them when their morphological features are indistinct. Moreover, eDNA sampling is non-invasive, and from the safety point of view, it is less labour intensive, employs less hazardous materials, and reduces health and safety exposure. Processing eDNA samples can be implemented using existing laboratory workflows and applied to a broad range of taxa without any special taxonomic expertise. Although there are some biological data that cannot be easily inferred from eDNA data, such as biomass, age, or health, the recent attempts to obtain at least some of these data from eDNA or eRNA are promising.

The consensus of this guidance is that the eDNA-based methods have the potential to be fully integrated into biomonitoring of energy project activities. The extent of their application depends on monitoring objectives as well as habitats and target taxa. Environmental DNA-based methods can, in some cases, be used as stand-alone tools for the detection of particular species (invasive, endangered, or pollution indicators). They could also be applied to assess environmental impacts and ecological status, although further validation of currently available eDNA-based metrics might be necessary. A combination of eDNA and conventional methods (such as imagery and/or acoustics) is appropriate for many applications. eDNA analyses provide a different type of information that is often complementary to what conventional biomonitoring can provide. The use of multiple methods, including eDNA-based techniques, will make future biomonitoring more powerful and comprehensive, which will help inform business decisions and aid in sustainable development within the energy industry.

The table below summarises where eDNA and conventional methods can currently be used for various habitats and target communities, at each major phase of operation for the energy industry. **Bolded** words indicate the most frequently used method. CA = Community analysis; SD = Species Detection

Habitat	Target community	eDNA sampling methods	Conventional methods	eDNA application						
				Recommended use	Exploration	Development (Baselining)	Production	End of Life		
								Decommissioning	Remediation	Restoration
Marine	Benthic sediment	Sediment sample; Macrofauna sample	Macrofauna sample; Benthic imagery; Active acoustics	CA with further validation	✓	✓	✓		✓	✓
	Benthic hard substrate	Surface scrape; Surface swabs; Surface suction	Surface scrape; Benthic imagery; Active acoustics	CA in combination	✓	✓	✓	✓	✓	✓
	Fish	Water filter	Netting; Benthic and aerial imagery; Active acoustics	CA & SD Stand-alone	✓	✓	✓		✓	✓
	Broad vertebrates	Water filter	Visual observations; Passive acoustics; Aerial imagery	CA & SD in combination		✓			✓	✓
	Zooplankton & Phytoplankton	Water filter; Bulk net sample	Water sampling; Netting; Active acoustics; Aerial imagery	CA in combination SD stand-alone		✓			✓	✓
Freshwater	Fish	Water filter	Netting; Electrofishing	CA in combination SD stand-alone	✓	✓	✓	✓	✓	✓
	Aquatic Invertebrates	Water filter Bulk sampling	Kicknet; Traps	CA with further validation SD stand-alone	✓	✓	✓		✓	✓
	Phytobenthos	Biofilm sampling	Biofilm sampling;	CA with further validation	✓	✓	✓		✓	✓
Terrestrial	Broad vertebrates	Water filter Air DNA; Surface swabs; iDNA	Visual and auditory; Camera traps; Vertebrate traps; Artificial cover	CA & SD in combination		✓		✓	✓	✓
	Invertebrates	Invertebrate traps; Air DNA; Surface swabs	Invertebrate traps	CA in combination		✓		✓	✓	✓
	Soil	Soil sampling	Soil sampling	CA with further validation	✓	✓	✓		✓	✓

Table of Contents

Disclaimer.....	i
Revision history.....	ii
Executive Summary.....	iii
Table of Contents.....	v
1 Introduction and Overview.....	1
1.1 Background.....	1
1.2 Scope and Objectives of this Guidance.....	1
2 Overview of Conventional Biological Monitoring Methods.....	2
2.1 Applications.....	2
2.1.1 Marine.....	2
2.1.2 Freshwater.....	4
2.1.3 Terrestrial.....	6
2.1.4 Remote Sensing and Other Survey Methods.....	8
2.2 Challenges and Limitations of Conventional Methods.....	9
3 Overview of DNA-based Methods.....	11
3.1 General Outline.....	11
3.2 Applications.....	13
3.2.1 Marine.....	14
3.2.2 Freshwater.....	17
3.2.3 Terrestrial.....	18
3.3 Summary of Challenges and Limitations of eDNA-Based Methods.....	19
4 Comparison of Conventional and eDNA Methods.....	20
4.1 General Outline.....	20
4.2 Marine.....	21
4.2.1 Vertebrates.....	21
4.2.2 Benthic Community.....	22
4.2.3 Zoo- and Phytoplankton.....	23
4.3 Freshwater.....	23
4.3.1 Fish.....	23
4.3.2 Aquatic Invertebrates.....	24
4.3.3 Phytobenthos (Diatoms).....	25
4.4 Terrestrial.....	26
4.4.1 Vertebrates.....	26
4.4.2 Terrestrial Invertebrates.....	27
4.5 Summary of Method Comparison.....	28
4.5.1 Fieldwork (Ease of Use, Sampling Effort, Cost, Safety).....	28
4.5.2 eDNA Sample Analysis (Processing Effort, Data Obtained, Reference Sequence Library, Turnaround Time).....	29
4.5.3 Output Data.....	29
4.6 Summary of Advantages of eDNA-Based Methods vs Conventional Methods.....	30

4.7	Recommendations	31
5	Application of eDNA-based Surveys in the Energy Industry	32
5.1	Review of Existing Literature	32
5.2	Conventional vs eDNA-Based Monitoring Using Offshore Platforms as a Model	34
5.3	Prospective eDNA Applications to Monitor Energy Project Activities	35
5.3.1	Baseline Surveys	37
5.3.2	Operations Phase	37
5.3.3	Oil Spills and Remediation	38
5.3.4	Decommissioning	38
5.3.5	Restoration	39
6	Elements to Consider When Planning eDNA-Based Surveys	39
6.1	Suitability	40
6.2	Feasibility	40
6.3	Reliability	41
6.4	Acceptance	41
6.5	Economic advantages	42
6.6	Added value	42
6.7	SWOT Analysis	43
6.8	Decision Tree for Selecting eDNA-Based Methods for Biomonitoring	44
7	Future Directions of eDNA Methodologies	45
7.1	General	45
7.2	Sampling Techniques, Equipment, and Surveying	45
7.2.1	Automated sampling	45
7.2.2	In-Field Analysis	45
7.2.3	Other Substrates for DNA Sampling	46
7.3	Laboratory Work and Analysis	46
7.3.1	Technological Innovations	46
7.3.2	Taxonomic Reference Database	46
7.3.3	Species Detection	47
7.3.4	Abundance Measures	47
7.3.5	Consistency of Results	47
7.4	Metrics of Ecosystem Health	47
7.5	Standardisation	48
7.6	Summary	49
	Glossary	50
	References	53

1 Introduction and Overview

1.1 Background

The energy industry undertakes a variety of ecological assessments, frequently as a regulatory requirement for proposed, existing or decommissioned assets. Conventional biodiversity survey methods, which usually have standardised protocols based on direct observations or collection of organisms, are broadly accepted by industry and regulators. However, these conventional methods can be labour-intensive, expensive, limited in taxonomic scope and require experts with taxonomic training.

Environmental DNA-based methods can complement, match and in many cases outperform conventional biodiversity survey methods (Keck, Blackman, et al. 2022; Valentini et al. 2016; Deiner et al. 2017). Their other advantages include reduced cost and survey effort, and increased taxonomic resolution. A recently published literature review identified that eDNA performed better than conventional methods in the majority of 194 comparative studies in terms of sensitivity (ability to detect species at low abundance), overall species richness detected and cost-effectiveness (Fediajevaite et al. 2021). The ease of field sampling for eDNA and its sensitivity has already made it a useful tool for species detection by professionals and citizens (Biggs et al. 2015). Environmental DNA-based methods can generate measures of ecosystem health (Pawlowski et al. 2018; Cordier et al. 2020). However, the current limitations of eDNA analysis in being able to quantify abundance and discrepancies between molecular and morphological identification can impede the uptake and acceptance by regulators of eDNA-based methods for biodiversity surveys.

The application of eDNA-based methods to the energy industry is still in a relatively early stage. Of the studies that have been conducted, their results show the immense potential of using DNA as an alternative to conventional monitoring. The detection of marine mammals during baseline assessment, the assessment of environmental impacts on sediment-dwelling fauna during operational phase, or the detection of invasive non-native species (INNS) during the decommissioning phase are just a few examples of potential applications for eDNA-based monitoring across different phases of energy project activity.

1.2 Scope and Objectives of this Guidance

This document is intended to provide environmental practitioners in the energy industry with guidance on when and where to apply eDNA methods. Recommendations within this guidance are based on a comprehensive literature review of conventional and eDNA-based biodiversity monitoring methods, with the aim to guide the selection of the most suitable methods for a given application, research objectives, study taxa and habitat. The document provides a critical overview of current applications of eDNA-based methods and, for a range of industrial applications, evaluates where these methods:

- can be used as stand-alone tools to obtain scientifically defensible data outputs.
- should be used in combination with conventional methods to generate the information required.
- are technically feasible but further testing and validation is required for industrial application.
- are not appropriate for use.

This report focusses primarily on the topics of field sampling for eDNA, data processing, and analysis as they relate to conventional sampling methods. Aspects including the regulatory acceptance, sample preservation, sample preparation and laboratory workflows, and data analysis are addressed where pertinent, but are covered to a greater extent in IOGP JIP34 chapters 2, 3 and 4.

2 Overview of Conventional Biological Monitoring Methods

2.1 Applications

A range of biological data are collected and analysed using conventional methods across the energy sector. These can include:

- species presence (and to a lesser degree, absence), temporal and spatial distribution
- species richness and evenness (alpha diversity)
- community composition and structure (beta diversity)
- abundance and biomass (absolute and relative; also using coverage as an abundance indicator)
- site condition/health (frequently based on community composition or indicator species presence/absence)
- physiological response to pollution (mussels and fish tissue analysis commonly used in energy industry)
- behaviour (less frequently recorded for use in the energy industry, but can influence other biological data metrics)

For the purposes of this document, we have focused on biological data that are commonly used in biomonitoring associated with energy industry operations that could potentially be obtained using eDNA-based methods. Conventional methods used for their collection are described below for marine, terrestrial and freshwater environments (Table 2.1).

Table 2.1 – Overview of Conventional Biological Monitoring Methods

Marine	Freshwater	Terrestrial	Others
Aerial imagery	Netting	Visual and auditory	Remote sensing
Visual observations	Electrofishing	Passive acoustics	Acoustic tagging
Passive & active acoustics	Kick net	Camera (infrared & thermal) traps	Crustacean trap
Underwater imagery	Sediment sampling	Vertebrate traps	Tissue sampling
Trawls/netting	Biofilm sampling	Artificial cover	Citizen science
Sediment sampling		Invertebrate traps	Box surveying
Plankton sampling		Soil sampling	Hand searches
		Microbial communities	

2.1.1 Marine

The main taxa targeted in marine environments are marine mammals, reptiles, seabirds, fish, benthic invertebrates and zooplankton. Aerial and vessel observer surveys are used for marine mammals, reptiles and seabirds, with passive acoustic monitoring also being applied to marine mammals. Fish data are most frequently derived from video and acoustic imagery and occasionally trawls. Benthic communities (those living at the bottom of the water body) are monitored either using video imagery targeting fish and sessile fauna or using either sediment grabs or cores to study benthic macro- and meiofauna, with active acoustics (sidescan sonar, MBES) used for identifying biotopes. Planktonic biota (those living in the water column) including zooplankton (ichthyoplankton and invertebrate plankton) and phytoplankton are monitored using trawled plankton nets and filtered water samples.

Acoustic and satellite methods can also be used for considering the distribution of phytoplankton.

Aerial surveys, Visual Observation and Passive Acoustic Monitoring. Marine vertebrates, such as mammals, seabirds, reptiles and elasmobranchs (cartilaginous fishes like sharks, rays, and skates), often include protected species, and monitoring them is required by current regulations. Conventional methods include aerial surveys, visual observations and passive acoustic monitoring to detect the target species in areas of energy project activity.

Dedicated aerial surveys, either via manned aircraft or unmanned drone, are frequently done prior to environmental baselining or even geophysical surveying. Aerial surveys typically use high-resolution imagery to detect species of interest, often combined with machine learning tools to reduce the cost of manual video analysis. However, this method is dependent on the target taxa being visible in the imagery which is affected by multiple environmental factors such as cloud cover and water turbidity.

Real-time visual observations are often required at fixed intervals during offshore operations because protected species such as marine mammals can be impacted by the noise from operations (such as geophysical surveys, pile-driving or drilling operations). Observations are often conducted from a survey vessel or platform. This requires constant observations throughout vessel deployment/offshore activity, with several marine mammal observers (MMO; UK) or protected species officers (PSO; USA) working in shifts. Depending on the size and scope of the project this type of monitoring can require large field teams. Observations and vessel activity are recorded and submitted to the relevant national authority, with data quality assurance/quality control (QA/QC) usually being carried out on land following project completion. Visual observations are generally limited by distance, sea state, and time spent on the surface by targeted species.

Simultaneously to visual observations, sound can be monitored using passive acoustic monitoring devices (PAMs). Detections from PAMs of marine mammal sounds can be carried out on board a vessel in real time or deployed on moorings for set periods of time. However, lengthy data analysis is frequently required to first pick out an acoustic signal and then to identify the species (Palmer et al. 2019).

Underwater imagery. Data extracted from video imagery are commonly used for investigating fish and sessile benthic fauna communities. Video imagery allows for the identification of species and the subsequent assessment of species diversity. It can also be used to count protected benthic species and measure the percentage cover and area of protected habitats (e.g. corals, sponge cover). Laser pointers or a fixed frame are commonly used to measure seabed features. Baited remote underwater video (BRUV) can be used for some mobile fauna, such as elasmobranchs (Boussarie et al. 2018).

The deployment of underwater video cameras requires trained personnel to oversee their use and depends on water turbidity and weather conditions. Obtaining video imagery results requires a high amount of effort, and has high associated field sampling costs, which can include the costs of equipment rental and deployment. The turnaround time for results of the video imagery analysis can be weeks from the date of sampling, although there is potential to reduce the human effort in analysis as machine learning technologies advance (Montereale Gavazzi et al. 2021). Video imagery remains a relatively expensive and time-consuming method. Moreover, imagery data may fail to detect cryptic taxa that are camouflaged, buried within the sediment or hidden in crevices.

Trawling. Fish and mobile pelagic and benthic macrofauna data are occasionally collected via midwater or bottom trawling. Such data are valuable because they can directly assess

the abundance and age of target species, however, the method is highly invasive and bottom trawling can have negative impacts on the benthic community (Jac et al. 2021) and surrounding ecosystem. The use of trawls can also entangle with cables and physical infrastructure. For these reasons, trawling is not a recommended method for surveying offshore energy project activity.

Sediment sampling. Sediment samples from benthic grabs or cores are commonly used for the assessment of benthic macro- and meiofauna. Benthic macrofauna-based indices such as AMBI (AZTI Marine Benthic Index, Borja, Franco, and Pérez 2000), are currently used for the assessment of impacts associated with offshore platforms activity. There are also indices that are tailored for particular groups of benthic meiofauna (e.g. The Maturity Index, Bongers 1990; foram-AMBI, Alve et al. 2016). All these indices are calculated based on a list of species assigned to ecological categories and their abundance. A benefit of sediment sampling is that useful assessments like the particle size distribution (PSD) and physicochemical variables (heavy metals, hydrocarbons and organic carbon) can also be obtained from the same sediment samples. A combination of imagery data, benthic macrofauna and PSD can be used to assign a biotope, such as when using the EUNIS habitat classification system (<https://mhc.jncc.gov.uk/>). The resulting habitat classification can then be used to assess environmental change and impact.

Collecting benthic macro- and meiofauna data requires substantial effort in the field, requiring sampling equipment deployment and retrieval, on-board sieving, and preservation in potentially hazardous solutions such as formaldehyde or industrial denatured alcohol. The time required for a laboratory to analyse samples from a project can be many weeks to months, with turnaround times often dependent upon the increasingly limited availability of taxonomists and sample complexity. Industry standards such as the NMBAQC (NE Atlantic Marine Biological Analytical Quality Control) give clear guidance on the taxonomic identification, procedures for reference collections and sample reanalyses (Cooper and Rees 2002). Nevertheless, such standards are usually limited to a certain biogeographic area or specific habitats.

Plankton sampling. Planktonic data used for biomonitoring consists mainly of marine zooplankton, which includes a range of invertebrate taxa (mostly copepods and cnidarians) and ichthyoplankton. Marine phytoplankton, which consist of various groups of algae, are less commonly targeted. Plankton samples are obtained from water either using plankton nets or by filtering water. They can be fixed in buffered formalin or with Lugol's solution.

The main challenge of analysing planktonic data is the taxonomic identification of typically small sized and often inconspicuous morphotypes. The automated analysis of plankton with imagery tools, such as the ZooCAM In-Flow imaging system allows for fast onboard counting, sizing and classification (Colas et al. 2018), so is becoming a more prevalent method to assess these communities, though other systems have been developed for the analysis microscopic plankton (Pollina et al. 2022). Despite these advances, the ability for such systems to accurately identify species is not optimal.

2.1.2 Freshwater

Fish, benthic invertebrates, and benthic diatoms are the main biological quality elements to be assessed during the monitoring of water bodies (e.g. EU WFD, Directive 2000/60/EC). Freshwater monitoring programs sometimes also include macrophytes and phytoplankton, but data on these taxa are less commonly used and will not be discussed here.

Netting and electrofishing. Freshwater fish populations are commonly monitored through netting or electrofishing. A range of nets can be used to survey fish, with the most popular being gill netting, seine netting and fyke netting. Fish captured through netting can be

identified, measured, sometimes sexed and any other relevant biological information taken (e.g. tissue samples). Several different netting methods can be used to ensure representative biological sampling, which may still reveal limited diversity (Hallam et al. 2021). This is in part due to the different conditions under which the netting techniques are effective and associated biases such as mesh size selectivity. This can result in species of interest not being assessed by netting. Logistically, a high sampling effort is required, especially in larger rivers and reservoirs, where entanglement of nets and trawls can be problematic.

Another widely used method for collecting fish data is electrofishing. The method consists of creating an electric field in a small area of water to immobilize fish, which then float to the surface and can be collected. This is a relatively comprehensive form of fish sampling, as it is less selective than most nets. The upstream wading electrofishing is used in streams and rivers, while a boating method is used in larger waterbodies that are too deep or wide to wade effectively. Once the sampling is completed, the crew records data on the catch, identifying each fish to species level, recording weights, and often noting any signs of external diseases or parasites. Fish are released back into the water once the data is recorded.

Both netting and electrofishing are invasive methods that can be harmful to fish. They also present a high risk from a health and safety perspective and require trained personnel. As each method is relatively selective in the capture of species not all species are surveyed equally. A risk of fish mortality can be minimised through careful equipment design and protocols which allows for catch and release. However, fish may not always be released unharmed due to unforeseen conditions or events (D. E. Snyder 2003).

Aquatic invertebrates sampling. Several groups of aquatic invertebrates are used as indicators of water and sediment quality and ecosystem health. Most commonly used are EPT insects representing the orders Ephemeroptera (mayfly), Plecoptera (stonefly) and Trichoptera (caddisfly), which are known to be particularly sensitive to water pollution. Another group of aquatic insects widely used as bioindicators of water quality in lakes are chironomids (non-biting midges) (Kranzfelder et al. 2015). Biological quality of sediment in rivers and lakes is commonly assessed using oligochaetes (Lafont et al. 2012) or nematodes (Höss et al. 2011).

Kick net sampling is the most popular method to collect aquatic insects and other invertebrates from rivers and streams. It is carried out by disturbing the river or stream bottom by kicking or shuffling and collecting organisms in a net downstream. Sweep net sampling is conducted in ponds and lakes to capture different taxa that inhabit aquatic vegetation. Sediment samples for oligochaetes and nematodes are collected using different types of nets, grabs or piston drills. The sediments are sieved and fixed with formaldehyde, and the specimens are sorted and morphologically identified under microscope.

To ensure that the sampling is representative of the site, it is important to sample different microhabitats (e.g. slow water, weeds, tree roots) and sampling effort can be allocated based on microhabitat prevalence. Identification to family level and taxon counts can then be carried out in the field and assigned an abundance category. The identification to species level is carried out in a laboratory under a microscope and usually necessitates an experienced taxonomist. Difficulties associated with the identification based on morphological features are the main factor limiting the use of aquatic invertebrates for ecological diagnostics.

Biofilm sampling for benthic diatoms. Monitoring water quality in rivers and streams using benthic diatoms (phytobenthos) is prescribed in Europe and recommended in other countries. Diatoms are effective for assessing water quality due to their rich diversity and

rapid growth, which enables them to respond quickly to changes in environmental conditions caused by natural or anthropogenic impacts. Since certain diatom species have preferences for different water quality conditions (nutrients, salinity, pH, heavy metals, temperature, and flow) the analysis of species composition in a sample can be used as a biological indicator of organic pollution, eutrophication, acidification and metal pollution. Among various groups of diatoms, those living as epilithic phytobenthos in rivers and streams are commonly used as bioindicators (Rimet and Bouchez 2012).

Sampling for diatoms is usually carried out by scrubbing the biofilm from upper surfaces of stones, placed in a sampling tray with stream water (Taylor et al. 2007; Gonzalo and de los Reyes Fernandez 2012). The samples are preserved in Lugol's iodine or ethanol and processed at a laboratory using a range of hazardous substances (Taylor et al. 2007; M. G. Kelly et al. 1998). Slides are prepared for microscopy, and the diatom valves are counted to measure the relative abundance of different species. The identification of freshwater diatoms requires expert taxonomic knowledge, but intraspecific variability can be very high, and the morphological differences between species can be so subtle that even trained taxonomists may come to different conclusions.

2.1.3 Terrestrial

Terrestrial biodiversity surveys typically combine a range of survey methods targeting different taxonomic groups. Methods include (among others) visual surveys for field signs of terrestrial vertebrates, the use of camera traps for large mammals and other vertebrates, mist netting for birds, mist netting and harp trapping for bats, acoustic surveys for amphibians, birds and bats, a variety of trapping approaches for small mammals, reptiles and invertebrates, and detailed plot surveys for vegetation. There is increasing use of remote sensing for initial assessments of habitat suitability and vegetation, but this is limited in its ability to deliver species specific data.

Visual and auditory surveys. Visual and auditory field surveys such as walkovers, transects and point counts are commonly used for species surveys in terrestrial habitats. The surveys are carried out by trained ecologists and can encompass a broad range of taxa including mammals, birds, amphibians, reptiles, and insects. Typically, surveys involve searching an area for field signs indicating the presence of the target species or directly observing the species. In the case of bird surveys, these are usually based on the identification of bird song rather than visual observations. Visual surveys may also include determining habitat suitability for the species of interest. Surveys are carried out at different times of day or night to capture the activity of different species groups and targeting the optimum season or spread across multiple seasons. They can be combined with other survey methods such as hand or dip netting, rock-turning for amphibians or searching buildings and trees for bat roosting features.

Acoustic surveys. Acoustic monitoring is widely used for bats but also for other terrestrial vertebrates (see Sugai et al. 2019 for a review). Acoustic surveying can capture more aerial insectivorous bat species than any other combination of sampling methods (Appel et al. 2022). Acoustic surveys may involve the use of handheld detectors during bat emergence surveys and activity transects, or static detectors may be deployed to record passively. These passive acoustic recorders do not require the presence of the researcher for the whole period and can be programmed to operate on a species-targeted schedule. Static detectors are deployed in different habitats within the survey area, positioning the microphones to ensure the optimal likelihood of detection. Typically, these are deployed for at least five nights in a row, often over a period over several months or across several seasons (Collins, 2023). Recordings are retrieved, then processed and analysed using call analysis software to identify calls to genus or species level where possible. Due to large

regional variations in the availability of audio reference libraries, the ability to identify to the species rank may be limited.

Camera trapping. Monitoring of mammals, reptiles and ground-dwelling birds can be carried out non-invasively using camera traps. These are widely used in wildlife monitoring projects worldwide - typically using remotely activated devices that capture images or video when a motion and/or heat sensor is triggered. The majority of studies using camera traps focus on mammals, with considerably fewer studies on birds and very few studies targeting reptiles or amphibians (Burton et al. 2015). They are subject to a range of detection biases, such as animals not entering the detection zone, the habitat characteristics (e.g. dense vegetation), attraction to a camera or even camera trap positioning and settings. These can be overcome to some degree by statistical modelling such as occupancy modelling (Wevers et al. 2021). As with all imagery, the camera traps images are reliant on human identification and counting of species by trained taxonomists, which is time consuming, although automated identification is increasingly being used (Schneider et al. 2020).

Traps for capturing mammals, reptiles and amphibians. Small mammals, reptiles and amphibians can be captured using a variety of traps. Ground-dwelling small mammals are commonly sampled using a baited box with a trapdoor, with modified designs depending on the size of target species (Torre et al. 2016). Pitfall traps using buckets placed in a line with drift fences can outperform the detection rates of other trapping methods for some species (Thompson and Thompson 2007; Mena et al. 2021). Trapped mammals can be tagged, allowing for the number of different individuals trapped to be assessed, and the population size estimated (Slade and Blair 2000). However, these methods have been found to have a lower detection rate for multiple species than camera traps (M. L. Thomas et al. 2020).

Artificial Cover Objects (ACOs) are regularly used for surveying reptiles, whereby ACOs create favourable refugia and can be accessed to count species and number of individuals on a frequent basis (e.g. Wilson, Mulvey, and Clark 2007).

Mist nets and harp traps are regarded as one of the most effective methods of sampling flying vertebrates (Trevelin et al. 2017). When targeting bats, mist nets are deployed near roost entrances, in areas with vegetation, or water where bats fly low (Ferreira et al. 2021) and in areas where they are more likely to be caught such as along commuting routes or at swarming sites. In addition, harp traps are widely used to capture bats. Thicker mist nets are usually used to capture birds in low visibility environments to complement visual and auditory methods (Marques et al. 2013). Birds in mist nets have been found to have a low (<1%) risk of injury and mortality (Spotswood et al. 2012).

Traps for insects and other arthropods. Insects with different modes of life are sampled using a variety of trapping methods and subsequently identified in a laboratory. As discussed below, the same sampling methods can also be adapted for DNA-based identification. These almost always kill the specimens sampled regardless of whether they are processed using DNA-based or morphological identification methods, which can be problematic when considering protected taxa. It can be illegal to trap if there are protected invertebrates in the area (e.g. rusty patched bumble bee in the USA; Franz 2020).

Flying insects are commonly collected using Malaise traps or pan traps. The Malaise trap is placed in a natural flyway, where hundreds of insects can be directed to the apex of the trap where they fall into a collection bottle of preservative solution. The trap can be deployed indefinitely, with only the bottle replaced periodically (Kirse et al. 2021), making it a very low maintenance and low-cost sampling technique. As it can be deployed in a highly standardised way, the Malaise trap can be used for biomonitoring programs that seek to track long-term trends, including the impacts of development or restoration activities on species diversity. Pan traps use containers that are partly filled with soapy water, which

breaks the surface tension of the water causing insects to fall in, rather than stay on top of the water. The water is usually dumped in a net and transferred to a preserving agent (usually ethanol) (Anderson et al. 2013). Other methods for sampling flying insects include: flight intercept and window traps, light traps, sweep nets for Hemiptera and Orthoptera and aerial nets for Lepidoptera and Odonata, and sticky traps primarily for agricultural pest monitoring (Böckmann et al. 2021).

Ground surface-dwelling arthropods (particularly beetles and spiders) are commonly collected using pitfall traps, which are containers sunk in the soil and filled with a preservative solution such as a mixture of water and ethylene glycol. This allows for high field replication due to the low handling time. Features such as funnels and guidance barriers can also be included to increase capture efficiency. However, there is no uniform method for sampling surface-dwelling arthropods, which can lead to variable results (Boetzi et al. 2018).

Collecting soil invertebrates. For smaller invertebrate fauna such as nematodes and microarthropods, soil sampling and subsequent sorting is the most commonly applied method, albeit rarely used at commercial scale due to the inherent time and effort. The sorting process involves a density-based separation (Schenk et al. 2020), and/or a filtration based sorting step. Sorting can be a labour-intensive and equipment-heavy process. When sample numbers are high, this requires substantial lab space. Invertebrate indicator species identified in soils can include earthworms, nematodes, mites and springtails. The metrics derived from these indicators generally incorporate abundance data of the target species group.

Monitoring microbial communities. Traditionally, soil bacteria and fungi have been assessed through culture-based methods (e.g. Janssen et al. 2002) or phospholipid fatty acid (PLFA) measurements (Frostegård, Tunlid, and Bååth 2011). A combination of culturing and molecular techniques are common approaches for identifying and monitoring microbial communities (Wani et al. 2022). Culture-based methods are generally considered selective and biased as a relatively small percentage of bacteria species can be cultured in the lab (e.g. F. O. P. Stefani et al. 2015). The varying concentrations of PFLAs are used as a rapidly obtained proxy for different microbial communities and biomass, particularly for the fungal to bacterial ratio (Frostegård, Tunlid, and Bååth 2011). However, PFLA analysis has largely been superseded by DNA-based methods for community characterisation.

2.1.4 Remote Sensing and Other Survey Methods

Currently, remote sensing is gaining attraction as a method for non-invasive surveys of aquatic and terrestrial environments over large areas using aircraft (including drones) or satellites, with minimal deployment of personnel. In terrestrial environments, remote sensing can produce high level 3D images of vegetation and measure plant biomass. In freshwater and marine environments, remote sensing is mainly used to estimate algae abundance (particularly chlorophyll concentration) and sediment load. It is an ideal tool to assess water extent and catchment hydrology, which are generally required and are highly relevant to water eDNA sampling (Carraro, Stauffer, and Altermatt 2021). Whilst habitat assessments derived from remote sensing are not measures of biodiversity, they can provide metrics such as Habitat Extent, Habitat Connectivity, NDVI (a ratio between the red and near infrared values, useful in understanding vegetation density and assessing changes in plant health), and soil type and health. The use of remote sensing as an environmental/ecological tool for habitat suitability and vegetation have been extensively discussed in a previous IOGP report (IOGP 2020).

Besides remote sensing, there are other biological survey methods that are frequently applied to biomonitoring with governmental targets or are led by conservation and academic research organisations. These other conventional methods include:

- Acoustic tagging of individuals
- Trapping (e.g. crustaceans, decapods)
- Tissue sampling for monitoring health or population genetics
- Active searches for terrestrial invertebrates (under microhabitats such as dead logs, under stones or plants associated with certain invertebrates)
- Citizen science (visual) observations
- Bird nest, bat and small rodent box surveys
- Feeder box surveys
- DNA analysis of faecal samples for species identification

Some of these methods, such as tissue sampling are used when monitoring energy industry effects on the potential accumulation of pollutants in wild caught fish or blue mussels. However, as these methods cannot be replaced or complemented by eDNA, they are not presented here.

2.2 Challenges and Limitations of Conventional Methods

The main challenges and limitations of conventional biomonitoring methods are summarized below:

- **High sampling costs:** conventional survey techniques are labour intensive and sometimes rely on complex logistics (especially in marine environments).
- **Slow sample processing and lengthy data analysis:** specimen sorting and morphological species identification is usually time-consuming, greatly increasing the period from sampling to data analysis. In some cases, such as video imagery and acoustic data, the turnaround time for data analysis can also be very long.
- **Low frequency of sampling:** the variations between samples can be very high, but repeated sampling is cost-prohibited
- **Limits of visual observations:** the accuracy of observations can be limited by environmental factors (weather conditions, water turbidity, etc.) and the behaviour of species (e.g. those buried in the sediments or hidden in crevices, etc).
- **Limits of morphological identification:** using morphological characters to identify species can often be challenging, and for many taxa, an identification to family or genus level is accepted. In general, morphological identification of small and inconspicuous morphotypes, including immature stages of life cycles requires highly skilled experts. Additionally, some common taxa comprise a multitude of cryptic species that can only be distinguished by genetic data.
- **Lack of taxonomic expertise:** the accurate identification of sampled organisms requires good knowledge of morphological taxonomy, yet the number of experienced taxonomists is rapidly decreasing, and the training of new taxonomists is rare.
- **Limited range of taxa sampled:** conventional methods are most often based on large size mega- or macrofauna that are relatively easy to observe and identify. Taxonomic groups with small body sizes or single-cell organisms (e.g. meiofauna and microbiota) are generally ignored, even if they are highly sensitive to environmental impacts.
- **Invasiveness or destructiveness of sampling:** Fishing or trapping of vertebrates, as well as collecting bulk invertebrate samples, may have negative effects on

populations of protected and endangered species. Some sampling methods such as bottom trawling can destroy habitat.

- **Hazardous products:** the solutions used in the preservation of collected specimens (formaldehyde, denatured alcohol) are potentially hazardous. This is also the case in substances used for processing biofilm samples (HCL, KMnO₄, hydrogen peroxide).
- **Pathogen risks:** conventional sampling procedures may increase the risk of pathogen transfer among sites on sampling equipment if not sterilised.

3 Overview of DNA-based Methods

3.1 General Outline

Environmental genomic methods can be applied to a range of habitats and systems, targeting diverse taxa, from bacteria to blue whales. This section reviews the main eDNA methods in use, particularly those successfully applied within the energy industry.

Environmental genomics methods using eDNA can be broadly divided into two groups: single species and multiple species applications. Each group of applications can be addressed with specific analysis methods. While this list is not comprehensive, a more detailed exploration and guidelines for environmental genomics workflows and their analyses can be found in IOGP JIP34 chapters 3 and 4.

- **Single species**
 - **Quantitative PCR (qPCR)**: A fluorescent signal is emitted as DNA gets amplified by PCR. This allows the DNA concentration to be quantified. This approach can be used to assess species presence/absence and its abundance, if a qPCR assay is properly validated.
 - **Digital PCR (dPCR)**: This method consists of partitioning the DNA into thousands of droplets (ddPCR) or microfluidic chambers and quantifying the number of positive amplifications in each partition. It is similar to qPCR in application but is more robust to inhibitors and offers greater quantitative precision of DNA concentration
- **Multiple species**
 - **Metabarcoding**: A standard DNA fragment (DNA barcode) from a sample containing many individuals is amplified through PCR and sequenced on a high-throughput sequencing platform.
 - **eDNA metabarcoding** refers to processing environmental samples (e.g. water, soil or sediment)
 - **Bulk metabarcoding** refers to processing a mixture of specimens sorted from environmental samples (e.g. by sieving, filtering or elutriation)
 - **Metagenomics**: the DNA from an environmental sample is sequenced directly without the PCR amplification of a specific gene region
 - **Metatranscriptomics**: the RNA from an environmental sample is directly sequenced to assess the “active” community.




Figure 1. Workflow of eDNA analysis for (4-6) metabarcoding and (4b) single-species detection.

The most popular eDNA-based methods employed in biomonitoring are metabarcoding and qPCR. Metabarcoding is commonly used when measuring diversity, whilst qPCR is mainly applied when targeting one or a few species (e.g. protected, commercially important, or invasive, non-native species). Metabarcoding and qPCR can both be considered cost-effective analyses where only a small number of species are of interest. However, for larger a number of species, metabarcoding is the preferred approach.

Validated qPCR assays can:

- be more sensitive for detecting individual species at low population levels, outperforming metabarcoding for a given taxon (Moss et al. 2022)
- have known limits of detection and the potential for quantification (see Thalinger et al. (2021) for a more in depth discussion of qPCR validation),
- have a greater propensity for facilitating in-field detection due to the lower complexity of steps and equipment involved (A. C. Thomas et al. 2020)

From a practical perspective, single species assays are cheaper, easy to standardise and provide a faster turnaround time, which can be essential when rapid action is recommended. However, the time and effort required to develop and validate qPCR assays is considerable compared to the use of standard metabarcoding methods. For a more detailed discussion of the merits metabarcoding and single species assays, see Bruce et al. (2021). The advantages of each method are summarized in Table 3.1 below.

Table 3.1 – Comparison of single species assays and metabarcoding

qPCR / dPCR	Metabarcoding
More sensitive for detection of single species	Many species can be detected simultaneously
Detection limits can be established	Detection can be improved through increase of sequencing depth
Target DNA can be quantified	Relative abundance can sometimes be inferred from number of reads (sequences)
Lower complexity and rapid analysis can facilitate application	Cost per sample can be reduced through multiplexing

When preparing to carry out an eDNA-based biodiversity survey, clear objectives must be set before selecting the eDNA approach, sampling methods, location and number of samples (Figure 2). These objectives in the context of energy project activity might be:

- baselining to describe biodiversity status and to identify spatial and seasonal variation in species occurrence
- monitoring species introduction to a location
- monitoring changes in community composition associated with an impact
- monitoring the recovery of initial state during decommission and restoration




Figure 2. Steps for consideration when carrying out eDNA-based biodiversity surveys

Once a project execution plan is determined, the eDNA biomonitoring workflow can be broken down into the following steps:

- Field sampling and preservation
- Sample preparation and DNA extraction
- DNA quality checks (concentration and/or purity)
- PCR and quality checks
- Library preparation and sequencing for metabarcoding
- Bioinformatics and reporting

The detailed description of steps in this workflow are provided in subsequent chapters in the IOGP JIP34 series of publications ‘Environmental Genomics Applications for Environmental Management Activities in the Oil and Gas Industry’ (Chapters 2 – 4).

3.2 Applications

This section addresses the common applications of eDNA-based methods used to sample different environments and taxonomic groups.

eDNA is commonly isolated from water, sediment or soil. Additionally, in the case of freshwater environments, biofilm DNA samples provide a good material for analysis of phytobenthos (mainly diatoms) and small-sized invertebrates. Recent studies have shown increasing evidence of the utility of airborne DNA, in particular to detect terrestrial vertebrates, flying insects and plants.

Bulk DNA from a pre-processed mixture of specimens can be used to generate a more comprehensive list of taxa that are not well represented (i.e. differ from traditional samples) in water or sediment eDNA samples (aquatic insects, marine benthic invertebrates, zooplankton). The material for bulk DNA analysis is generally collected in the same way as for morphological studies, except for extra steps in decontamination, preservative solutions, and the laboratory processing.

Below, we present the different types of samples used for DNA analyses for different environments and target taxa (Figure 3).




Figure 3. Potential applications of eDNA-based methods for different environments, sample types and target taxon groups.

3.2.1 Marine

DNA-based monitoring of marine biodiversity is mainly applied through the analysis of sediment and water samples. Sediment eDNA is generally used to assess changes in the benthic macro- and meiofauna community, while water eDNA tends to be used to assess a broad range of vertebrates as well as planktonic taxa. Soft bottoms are preferentially sampled for eDNA, either through the analysis of sediment samples or bulk samples. eDNA analysis of hard bottoms is more challenging.

Soft bottom sediment samples from grabs or cores are commonly collected during conventional monitoring activities as well as during geotechnical surveys. These samples can be used for eDNA-based monitoring approaches, provided that the surface is undisturbed. Sediment is subsampled, frequently to a depth 2-3 cm, where most of epibenthic species are found. However, if the focus is on infauna, subsurface layers down to 10 cm should be subsampled. These subsamples can then be analysed separately or pooled and subsampled. For reviews of the methods of sediment sampling for eDNA and its preservation, see Pawlowski et al. (2022) and Wort et al. (2022).

Sediment eDNA samples can be used to monitor a wide range of taxonomic groups. Prokaryotes have been used to assess the functional composition before and after contamination events (Krolicka et al. 2014), to identify potential resilience, track site remediation (Wang et al. 2013), and to monitor restoration activities (e.g. Laroche et al. 2018). Microbial eukaryotes, meiofauna and traces of benthic macrofauna present in sediment eDNA samples have been extensively used for environmental assessment of

impacts associated with marine aquaculture (Pawlowski et al. 2014), and O&G activity (Lanzén, Mendibil, et al. 2021).

Vertebrate groups such as teleost fish and elasmobranchs can also be detected in sediment DNA, but this analysis typically recovers less diversity than water eDNA samples (Koziol et al. 2019). Furthermore, there is a high degree of temporal uncertainty regarding eDNA preservation in sediments as the DNA often has a longer residence time than in water (Kuwaie et al. 2020; Sakata et al. 2020). This means that baselining fish communities from sediment may not reflect the community at the time of sampling or show community changes associated with acute impacts.

Marine hard substrates are rarely sampled for eDNA due to difficulty in using grab or core samplers (but see Alexander et al. 2022; Alexander, Marnane, McDonald, et al. 2023). This may be resolved by a suction based sampling approach as proposed by Keeley et al. (2021) (Figure 4). The substrate type can influence the communities detected, hence any sampling must be representative of the local heterogeneity and be appropriate for the survey objectives (Koziol et al. 2019). For shallow or intertidal hard bottom samples, scrape or swab samples obtained by divers or ROVs can be used (Wangensteen, Cebrian, et al. 2018; Wangensteen, Palacin, et al. 2018; Alexander, Marnane, Elsdon, et al. 2023). For deep-sea sites, the bottom water can be used as a source of eDNA (Günther et al. 2018), although there is low taxonomic overlap between communities represented in benthic water and substrate samples (Laroche et al. 2020a).

Bulk samples of benthic macroinvertebrates can be used as an alternative of sediment sampling. In this case, sediment samples are sieved on-board, as performed in conventional benthic monitoring, and then the samples are preserved for subsequent DNA extraction and environmental genomic workflows. If the sediments contain a large amount of mineral debris (shells, coral skeletons) it is preferable to extract DNA from the preservative solution (i.e., ethanol) rather than from the homogenized material. The analysis of bulk DNA from the preservative enables identification of specimens using both morphology and DNA and provides data that are more similar to conventional methods compared to extracting eDNA from sediment. However, the taxonomic composition of bulk DNA samples are often different from conventional observations (Cewart et al. 2015; Loos and Nijland 2021), not to mention that smaller components of the benthic community (meiofauna and microbiota) are lost, reducing the range of community level data that can be obtained from eDNA analysis.

Bulk samples can also be used for zooplankton DNA analysis. In this case, plankton nets are used to collect specimens from surface waters. Samples of mixed specimens can be stored in ethanol or DNA can be extracted directly from the mixed specimens. However, as the zooplankton communities can fluctuate due to environmental and seasonal changes, weekly or monthly repeated sampling is recommended (Song et al. 2021).




Figure 4. Suction based substrate sampler for hard substrate. Image and publication credit to Keeley et al. (2021).

Seawater sampling design should consider that eDNA can be highly localised (both vertically and horizontally), particularly in low-energy, slow-moving environments and when the water column is stratified (Port et al. 2016; Yamamoto et al. 2017; Jeunen et al. 2020; Djurhuus et al. 2018; Ely et al. 2021). Samples from the surface, at the thermocline, and near the bottom can have different biological communities. Knowing oceanographic conditions is key to ensuring samples are taken at appropriate depths for interpretation of results. Niskin or GoFlo bottles are often used in an offshore setting (e.g. Closek et al. 2019), although nearshore, surface water samples can be taken using similar methods as sampling in freshwater (described below).

Autosamplers can be deployed for eDNA sampling at fixed intervals at a given location (Mynott 2019; Formel et al. 2021), or deployed on autonomous underwater vehicles (AUVs) (Hansen et al. 2020; Yamahara et al. 2019b; Truelove et al. 2022). An alternative is to use passive eDNA samplers which do not filter water (Bessey et al. 2021). These may be suitable for time-sensitive biological surveys, because they can collect eDNA over an extended duration if the adsorbent material is deployed appropriately. Research into passive sampling methods has developed rapidly in the past two years, with a wide variety of materials and protocols being assessed (Jeunen et al. 2022; Verdier et al. 2022; Kirtane et al. 2020).

Seawater eDNA samples are usually filtered and preserved with a buffer (e.g. Longmire's solution) and/or frozen once recovered on the vessel (Bruce et al. 2021). Mechanical filtration using a pump is recommended to reduce the time and effort processing and allowing a greater volume of water to be filtered. This is key for marine sampling where the water typically has a lower concentration of eDNA than freshwater. Sampling can be carried out by personnel with minimal training in most conditions. Furthermore, fewer personnel are required than with conventional sampling, resulting in less hours on deck and hence less risk to safety. For further detail on water eDNA sampling, refer to IOGP JIP34 Chapter 2

publication on 'Development of Industry Guidance on eDNA Sampling Standards and Guidelines'.

Seawater eDNA can be used to detect a wide range of pelagic taxa, from vertebrates to bacteria. While marine mammals and fish are successfully detected with eDNA, elasmobranchs, reptiles, and seabirds have been more problematic (although see Farrell et al. 2022; Boussarie et al. 2018; Liu et al. 2022). In the case of elasmobranchs and reptiles, this is thought to be due to their low relative abundance and the low concentration of eDNA shed by these organisms (Adams et al. 2019). In these cases, sampling a greater volume of water can increase the detection probability of these rare species (McClenaghan et al. 2020; Schabacker et al. 2020).

Among other organisms, bacteria are frequently analysed from marine water samples following oil spills and in the subsequent remediation phase (Brakstad et al. 2015; Dubinsky et al. 2013; Krolicka et al. 2019; Zhang et al. 2018), with some studies also applying metatranscriptomics to show the active bacterial community (Knapik et al. 2020). These studies consistently use a finer filter mesh (0.22 µm) to enhance the capture of these microorganisms.

3.2.2 Freshwater

DNA-based methods for freshwater biodiversity monitoring principally use a filtration kit and a standard sampling protocol. Bulk invertebrate samples as well as biofilm samples can be used to target selected groups of freshwater bioindicators (aquatic insects, oligochaetes, nematodes, diatoms) when investigating ecological status of water bodies.

Freshwater samples contain the DNA of organisms that spend part or all their lives in water as well as those from the surrounding areas, due to surface runoff of biological material and animals using it for drinking and bathing. This means that freshwater samples can be analysed for a broad range of aquatic, semi-aquatic and terrestrial species, allowing for rapid surveying of both freshwater and terrestrial environments (see also section 3.2.3.).

Freshwater eDNA is commonly used to assess the diversity of fish and amphibians. Many peer-reviewed articles report using eDNA to assess fish diversity or to detect particular species (Hänfling et al. 2016; Bradley et al. 2022). The utility of eDNA for amphibian monitoring is best illustrated by the example of a great crested newt assay used since 2014 in the UK that has broadly replaced habitat suitability surveys with a qPCR test (Biggs et al. 2014; Rees et al. 2014).

Freshwater eDNA has also been used successfully to detect aquatic molluscs (Blackman et al. 2020) and groups of insects used as bioindicators (e.g EPT; Brantschen et al. 2021). The water eDNA analysis of microbial communities (prokaryotes and eukaryotes) has been shown to have a broad range of applications, such as detecting pathogens and parasites (Bass et al. 2023), as well as the assessment of agricultural and human waste pollution impacts (Jin et al. 2018; Green et al. 2014; Malayil et al. 2020). More specifically, bacteria from water samples have also been analysed for assessing the impact of oil spills and hydrocarbon degradation (Irfan Ali Phulpoto 2021; Jurelevicius et al. 2013; Tiburcio et al. 2021).

During field sampling, water eDNA samples are commonly filtered *in situ*, through manual or mechanical pumping (A. C. Thomas et al. 2020; Bruce et al. 2021). In some cases, water samples are frozen to minimize DNA degradation and subsequently thawed and filtered in a lab, or water samples are kept cool for transport back to a lab for filtering. During the early adoption of eDNA, other methods of eDNA capture were also used, such as ethanol precipitation (Biggs et al. 2014) and centrifugation (Klymus et al. 2017). Both methods are currently considered as inefficient, as only a limited volume of water can be processed,

resulting in a low DNA yield (Bruce et al. 2021). Nevertheless, ethanol precipitation is still used in the UK for one of the few regulated eDNA survey methods (detection of great crested newts, Biggs et al. 2015).

The DNA in water typically degrades within hours or days, thus the eDNA analysis focused on macrofauna usually captures fine-scale local (horizontal and vertical) and temporal communities. The persistence of eDNA in water depends on a range of factors, including:

- Water transport and mixing (flow, stratification, wave action)
- Temperature
- Water chemistry (e.g. pH, salinity, conductivity)
- Microbial activity
- UV-B radiation
- Substrate (the underlying layer – in rivers, streams, oceans etc.)

Some of these factors can be addressed by sampling strategies or analyses that consider seasonality, waterbody size, type, depth, and substrate. In particular, for rivers and streams, currents can result in downstream transport of eDNA (Spence et al. 2021; Pont et al. 2018; Wacker et al. 2019). For enclosed ponds and lakes, eDNA can be highly localised due to lack of flow and wave action (Brys et al. 2021; J. Li et al. 2019; Shu et al. 2022). Similar to marine systems, vertical eDNA transport may be limited in deeper lakes with thermal stratification (Littlefair et al. 2020).

Bulk invertebrate samples are an alternative method for eDNA analysis of invertebrates living on the surface or in the sediments and are often used as bioindicators of environmental quality of freshwater bodies. Bulk DNA samples can generally be collected using the same procedures as the conventional methods outlined in section 2.3.1. with the principal differences being the preservative solution (usually ethanol), additional steps required to decontaminate equipment, and the method for sample preparation depending on the target taxa. The most common method for freshwater invertebrates collected using kick-nets is homogenisation of the bulk sample (e.g. Elbrecht et al. 2017; Pereira-da-Conceicao et al. 2020). However, eDNA extraction from the sample preservative has also been shown to provide satisfactory results (Zizka et al. 2019). Similar bulk eDNA protocols apply for aquatic oligochaetes and nematodes, yet in the case of oligochaetes the best results were obtained using high-throughput sequencing of sorted and genetically tagged specimens (Vivien et al. 2020).

Biofilm samples can be used as a source of eDNA for monitoring the ecological status of rivers and streams using diatoms. Biofilm sampling methodology for eDNA is the same as for conventional monitoring outlined in section 2.3.1, following standard protocols (NF EN 13946 – April 2014) as set out under the Water Framework Directive (WFD). A Technical Report from the European Committee for Standardisation (CEN/TR 17245, 2018) provides recommendations to maintain compatibility of eDNA samples. This can be obtained by preserving biofilm samples in 70% ethanol (e.g. Pérez-Burillo et al. 2020) or by using commercial DNA preservation buffers (Visco et al. 2015; Apothéoz-Perret-Gentil et al. 2017). Biofilms can also be passive samplers for freshwater fish and macroinvertebrates eDNA, although their capacity to detect rare species was rather limited (Rivera et al. 2021; 2023).

3.2.3 Terrestrial

Terrestrial taxa can be surveyed using eDNA-based methods in a variety of ways. The most common are the analysis of soil, water or bulk eDNA samples. There are also several alternative DNA-based approaches to monitor terrestrial communities that include:

- faeces (Guillerault et al. 2017)
- insect-derived DNA (Gogarten et al. 2020)
- airborne DNA (Lynggaard et al. 2022)
- snow (Franklin et al. 2019)
- floral swabs (Newton et al. 2023)
- rollers on hard substrates (Kyle et al. 2022; Allen et al. 2023; Valentin et al. 2020)

Soil eDNA can be used to assess the diversity of bacteria (Nacke et al. 2016), fungi (Danielsen et al. 2021), invertebrates (Bienert et al. 2012) and terrestrial mammals (Leempoel, Hebert, and Hadly 2020). Soil microbial communities are commonly used for terrestrial restoration studies (van der Heyde, Bunce, and Nevill 2022). Soil fungal and bacterial responses to environmental change (e.g. pollution) are often more rapid than changes in the plant communities, which makes them excellent proxies for monitoring shifts in ecosystems (Bahnmann et al. 2018).

Soil sampling is usually carried out using composite subsamples (combining several small soil samples) within a given area to reduce small-scale variability. Sampling depth must be kept constant between samples. Subsamples can be collected using a mini-corer, trowel/scoop or where the ground is particularly hard, a metal corer. Samples are frozen or preserved in a buffer, and small subsamples (1 – 10 g) are used for final DNA extraction (Bruce et al. 2021).

Water samples can be used to detect terrestrial vertebrates, which leave trace DNA when drinking, wading, swimming, urinating or defecating in water (Ushio et al. 2017; K. E. Williams et al. 2018; Harper et al. 2019). However, detection of terrestrial taxa can be sporadic as their contact with water is not constant. Consequently, their DNA is likely to be present at lower concentrations than aquatic or semi-aquatic taxa which may result in a lack of detection in some samples. Detection of terrestrial taxa is highly dependent on the level of sampling effort, sample replication, sample representativeness, and species either directly interacting with water and/or their DNA being transported from other waterbodies/via runoff. Furthermore, eDNA sampling from water is not often possible in arid or frozen conditions.

Bulk DNA samples derived from different types of traps are frequently used for DNA analysis of larger mobile terrestrial invertebrates (isopods, myriapods, insects). The bulk samples are collected using the same procedures as for conventional means (e.g. Piper et al. 2019; Oliverio et al. 2018). The differences for eDNA-based methods include using ethanol for preservation and the bulk samples are often homogenised (Oliverio et al. 2018). However direct DNA extraction from the preservation buffer is being used more often to allow for subsequent morphological identification (Drummond et al. 2015).

3.3 Summary of Challenges and Limitations of eDNA-Based Methods

The main challenges and limitations of eDNA-based methods are:

- **Persistence of eDNA** in water, sediment and soil depends on a range of environmental and biological factors, which should be taken into consideration.
- **eDNA disperses** in water over long distances making it difficult to determine exact location where species occurred.
- **Species detectability** in water eDNA (freshwater or marine) depends on DNA excretion (how much of DNA is shed by target species to the environment), degradation, and transport or diffusion (e.g. elasmobranchs and reptiles are difficult to detect due to their relative low abundance and low concentration of eDNA shed by these organisms).

- **Taxonomic resolution** of closely related species (ability to distinguish them) can be difficult with commonly used DNA barcodes.
- **DNA accumulation** over time in sediment and soil may impede temporal analysis of collected data.
- **Small volume of sediment or soil** (1-10 g) used for DNA extraction make it difficult to obtain a representative sample of larger organisms such as macroinvertebrates.
- **Hard bottoms** are difficult to sample for eDNA.
- **Collecting bulk DNA samples** can be labour intensive (no difference compared to conventional methods).
- **Bulk DNA** provides a reliable inventory of species, but their relative abundance is difficult to infer due to biomass and methodological biases.
- **Substrate availability** may be limited (e.g.no water for a site), resulting in sub-optimal eDNA methods being feasible for given taxa at a location

4 Comparison of Conventional and eDNA Methods

4.1 General Outline

This section compares conventional methods and eDNA-based methods for each major biological group in marine, terrestrial and freshwater habitats. This comparison is based on a literature review of 138 publications which can be found in Appendix A available as a separate excel document. These were reviewed and summarised in tables (appendices) detailing:

- Location and habitat
- Conventional sample type
- DNA sample type and target
- Whether more taxa were detected with eDNA-based methods vs. conventional methods
- Whether more useful community analyses were obtained from DNA samples
- Taxon overlap between sampling methods
- Correlation of abundance/biomass and DNA (read numbers, qPCR or dPCR outputs)
- Whether authors recommend eDNA-based methods as a replacement

Results were divided into broad taxonomic and ecological categories (biological groups), as in the previous sections. The following groups are considered:

Marine:

- Vertebrates (fish, marine mammals, reptiles; birds and elasmobranchs)
- Benthic invertebrates
- Zooplankton and phytoplankton

Freshwater:

- Fish
- Aquatic invertebrates (molluscs, arthropods)
- Phytobenthos (diatoms)

Terrestrial:

- Vertebrates, including semi-aquatic taxa such as amphibians
- Invertebrates

This section focuses on the 'target taxa' that were directly compared within the reviewed literature to provide concise and clear comparisons. This results in a more conservative

assessment of eDNA-based methods relative to conventional methods, as it does not consider the whole spectrum of taxa which can be detected using the same sample (e.g. other vertebrates where mammals are targeted). This comparison downplays one of the inherent advantages of eDNA metabarcoding of targeting a broad range of taxa and recovering non-target taxa from an analysis at the same time. For this report, taxon/taxa will be used for comparing number of OTUs or ASVs with the number of morphospecies.

4.2 Marine

4.2.1 Vertebrates

A total of 13 publications were reviewed that made comparisons in marine vertebrates. Eight papers focused on fish, two on aquatic reptiles, two on marine mammals and one broadly assessed vertebrates. The studies were done on different habitats, including coral reefs, kelp forests and subarctic waters. In 11 studies, the eDNA and conventional data were obtained simultaneously, with conventional data obtained through underwater visual surveys, video ROV, trawl catch, and surface visual observations. In two studies, the comparison was done based on previous observations. Most studies used eDNA metabarcoding, with only two studies using qPCR/dPCR for single species detection.

At a broad taxonomic level, the authors highlight a good congruence between eDNA and conventional methods, especially for fish and marine mammals. For example, 26/31 families of deep-water fish were found with both eDNA and trawling (Thomsen et al. 2016). However, at the species level, taxa identified through eDNA metabarcoding only partially overlapped with those found through trawling or visual surveys (Closek et al. 2019). In general, the eDNA methods detected a significantly greater number of species (up to three times as many) than were reported in visual surveys (Alexander et al. 2022; Lamy et al. 2021). This difference was particularly high in tropical waters, where more taxa were detected in eDNA samples than in underwater visual surveys. However, in tropical waters, fewer taxa were identified to species level (Polanco Fernández, Marques, et al. 2021; Valdivia-Carrillo et al. 2021), likely resulting from gaps in DNA barcoding reference libraries which will continue to improve over time. By comparison, the publicly available reference database for marine mammals and northern hemisphere fish species is relatively complete (Miya, Gotoh, and Sado 2020) making those eDNA studies much more effective.

Compared to fish and marine mammals, the effectiveness of eDNA was lower for aquatic reptiles, as indicated by the inconsistent detection of turtles, snakes and skinks in Australian estuaries (West et al. 2021). These species were semi-aquatic or terrestrial which could explain the difficulties in detecting them in water eDNA. Turtles and elasmobranchs may shed less DNA material than other taxa, leading to a lower concentration of eDNA, and therefore greater difficulty in detection (Closek et al. 2019). These limitations can be overcome through the use of targeted primers and including sequences with low counts for elasmobranchs (Boussarie et al. 2018), or use of qPCR assays targeting turtles (Farrell et al. 2022).

Environmental DNA detects a broader phylogenetic diversity and is more effective at detecting rare species known to be in the area but not observed visually within the study (Port et al. 2016; Polanco Fernández, Marques, et al. 2021). It also allows for the simultaneous identification of fish and marine mammals as part of a broad vertebrate assessment. eDNA outperforms diver counts, allowing for the observation of spatial patterns at fine scales with less effort (Lamy et al. 2021). Finally, eDNA can contribute additional data that can help practitioners understand and conserve threatened species such as population-level information in sea turtles (Farrell et al. 2022) and whales (Baker et al. 2018). However, authors agree the best approach to produce a comprehensive survey that includes the

broadest range of taxa is the combination of eDNA and visual surveys (e.g. Alexander et al. 2022).

4.2.2 Benthic Community

Twenty-four publications were reviewed to compare the efficacy of eDNA versus conventional methods for assessing benthic biota. Conventional benthic biomonitoring is often based on an assessment of macroinvertebrates (and less frequently meiofauna), obtained through grab or core sediment sampling. Some of the reviewed papers directly compared the results obtained through analysis of sediment DNA with morphological identification of macrofauna (Aylagas et al. 2018; Cowart et al. 2015; Steyaert et al. 2020; Klunder et al. 2020; Cahill et al. 2018). Other studies used the indices based on benthic macrofauna as a benchmark, against which the eDNA detections of macrofauna as well as other groups were compared. The authors tested whether DNA data provided the same ecological quality assessment as conventional benthic macrofauna surveys. In addition, some papers compared molecular and morphological data for meiofauna (foraminifera and nematodes).

The papers generally reported a low overlap in taxa between conventional and eDNA-based methods. Metabarcoding was found to detect only 36% and 50% of macrofauna species identified by morphological surveys in seagrass (Cowart et al. 2015) and estuarine habitats (Aylagas et al. 2018), respectively. Both methods were unable to identify the same set of species, even when the number of observed taxa were similar (Steyaert et al. 2020; Cahill et al. 2018). In most of the studies examined, sediment metabarcoding detected more taxa (OTUs/ASVs) than conventional methods. For example, the number of benthic taxa revealed by metabarcoding was on average three times higher in sediments near offshore gas platforms in the North Sea compared to conventional surveys (Klunder et al. 2020). This was attributed to the inclusion of broader groups and cryptic taxa in smaller size classes, which was particularly notable when comparing metabarcoding results from sieved and unsieved samples, where fewer taxa were identified in sieved samples (Steyaert et al. 2020).

The differences in taxonomic composition between metabarcoding and morphological methods might be explained by the small volume of sediments analysed for eDNA-based methods, subsequently resulting in a small fraction of the macrofauna community being detected (Aylagas et al. 2016; Pawlowski et al. 2022). Some papers also highlight the issue of a lack of coverage in reference database, which leaves numerous sequences unassigned (Cowart et al. 2015; Steyaert et al. 2020). Another reason may be primer bias that can impede detection of some common genera of nematodes (Rzeznik-Orignac et al. 2017). In the case of foraminifera specifically, the difference between molecular and morphological data was explained by the fact that conventional methods consider only shelled taxa, while eDNA analysis detects shelled and soft walled taxa (Frontalini et al. 2020).

Despite the differences in richness and taxonomic composition, the consensus from the papers indicates that metabarcoding data provide similar ecological quality status (EQS) as conventional benthic macrofauna surveys. This can be partly related to a correlation between biomass of dominant macrofauna indicators and their relative abundance in eDNA data (Lejzerowicz et al. 2015; Klunder et al. 2020). However, when macrofauna were not well represented in results from sediment DNA due to a lack database completion, a taxonomy-free approach has been proposed to overcome the absence. A supervised machine learning approach was introduced to predict the EQS from bacteria or microbial eukaryotes, using macrofauna indices as benchmark (Cordier et al. 2017; 2018; Frühe et al. 2020). In addition, a multi-trophic metabarcoding biotic index was created for monitoring benthic communities (Keeley, Wood, and Pochon 2018).

The consensus of the reviewed papers was that metabarcoding shows great promise for benthic monitoring of soft and hard bottoms. Although the use metabarcoding data comes with many challenges (reviewed in Duarte et al. 2021), several studies show that macrofauna-based indices can be efficiently replaced by metabarcoding analysis of other benthic taxa (meiofauna, protists, or bacteria). Depending on whether the aim of monitoring is to provide the inventory of soft-bottom community or to assess its ecological status, metabarcoding data can be used as a complement to, or replacement of, conventional methods. For hard-bottom fauna the combined approach of eDNA and video imagery is recommended to detect the most taxa, because any single method will miss some species.

4.2.3 Zoo- and Phytoplankton

A total of five publications were reviewed for pelagic non-vertebrate community comparisons. Three papers compared taxonomic composition of zooplankton community identified by microscopy to metabarcoding data obtained from seawater eDNA and/or bulk plankton samples collected using plankton trawls or a Continuous Plankton Recorder (CPR) (Deagle et al. 2018; Djurhuus et al. 2018; Suter et al. 2021). One paper estimates phytoplankton relative abundance compared to quantitative optical data (Pierella Karlusich, Pelletier, et al. 2022). The fifth paper analysed the efficiency of a qPCR assay for the detection of pelagic larvae of the invasive European green crab (Roux et al. 2020).

The eDNA-based zooplankton community studies consistently found a higher number of species compared to microscopic analyses, up to 1.6 times in the case of eDNA metabarcoding and even higher for bulk samples (Suter et al. 2021). Although significant differences in taxonomic composition were observed, all dominant taxa were detected by both molecular and microscopic analyses (Djurhuus et al. 2018; Kim et al. 2019). Regarding zooplankton quantitative data, the results of eDNA and bulk metabarcoding would require considerably more calibration to obtain relative abundance data (Deagle et al. 2018). However, it has been shown that biomass calculations for some zooplankton species can be derived from qPCR assays, where the gene copy number has been found to correlate with biomass (Jungbluth et al. 2022). Similarly, a robust metagenomic approach was proposed to estimate relative abundance of phytoplankton targeting a photosynthetic gene (Pierella Karlusich, Pelletier, et al. 2022).

The papers conclude that eDNA metabarcoding is a promising technique for plankton biomonitoring. However, they also highlight several challenges related to the use of eDNA-based methods, including cross-contamination issues while sampling, gaps in reference database, and biases in quantitative data (Deagle et al. 2018). Further refinement and standardisation seem necessary to make the methods more reliable (Suter et al. 2020). A combined approach of conventional (netting and more recently developed imagery approaches) and eDNA-based methods is recommended (Ibarbalz et al. 2019).

4.3 Freshwater

4.3.1 Fish

A total of 26 publications were reviewed that compared conventional and eDNA-based approaches to detecting freshwater fish communities. The papers report the eDNA data from different water bodies, including lakes, rivers, streams and ponds. Most articles characterize fish community, comparing taxonomic composition and relative abundance inferred from eDNA to data obtained using various conventional methods. Few studies focus on single species detection, which is important for conservation or management of invasions.

In most publications, more taxa were found using eDNA-based analyses than with conventional methods, with the exception of emptied/drained ponds, where essentially all fish were captured (Blabolil et al. 2022; Di Muri et al. 2020). One study using a combination

of conventional methods also detected a greater number of taxa than the eDNA-based survey alone (Nakagawa et al. 2018). Overall, eDNA appears to be the optimal method for assessing fish diversity, identifying a greater species richness with equal or lower sampling effort than conventional methods. This frequently led to more informative community analyses (Hayami et al. 2020; Hervé et al. 2022; Lawson Handley et al. 2019; J. Li et al. 2019; M. R. Snyder and Stepien 2020).

The reviewed literature indicated that a high proportion (60 - 100%) of fish species identified using conventional methods were also detected using eDNA surveys. In addition, eDNA-based surveys can have a lower risk of false negatives relative to conventional methods (Blabolil et al. 2022; Bradley et al. 2022). However, the identification to species rank using eDNA can be challenging due to incomplete reference databases, especially in the tropics (Doble et al. 2020). Another challenge is the lack of taxonomic resolution of common DNA barcodes between some closely related species.

Fish abundance and biomass data generally correlated with the number of sequence reads or qPCR results (e.g. Pont et al. 2022), but this relationship also varied depending on the species, environmental conditions and life history traits (see Appendix for more details). Very few publications considered eDNA-based approaches to be a complete replacement for conventional methods, particularly with regard to abundance (Di Muri et al. 2020; 2022; Boivin-Delisle et al. 2021), but considered eDNA-based sampling best used in combination with conventional methods. In this scenario, fish size and abundance data can be collected in accordance with regulations such as the Water Framework Directive, and the use of eDNA analysis can allow for an overall reduction in sampling effort and cost.

An important difference between eDNA-based and conventional freshwater fish surveys is that multiple conventional sampling methods are commonly used in parallel as each method is selective, meaning only part of the breadth of biodiversity can be detected (Hallam et al. 2021; Golpour et al. 2022; Pukk et al. 2021). By contrast, eDNA surveys frequently show a capacity for detecting a broad range of species, including elusive taxa that were previously undetected by conventional methods (Hallam et al. 2021).

4.3.2 Aquatic Invertebrates

A total of 16 publications were reviewed to compare eDNA-based methods and conventional methods used for assessment of freshwater invertebrates. The eDNA-based invertebrate community data were compared to kick-net sampling and other conventional methods used for the assessment of water and sediment quality. The targeted taxa included aquatic insects, molluscs and oligochaetes (the latter isolated from sediment and Surber samples). Some papers also used eDNA methods to detect invasive macroinvertebrate species (e.g. quagga mussels; Blackman et al. 2020).

Most of the studies used eDNA from filtered water, although in some cases bulk invertebrate samples were analysed. Bulk invertebrate samples often produced a higher diversity of target taxa (Harper et al. 2021b; Pereira-da-Conceicao et al. 2020), but were more localised in their representation compared to eDNA from water samples. This may reflect incompleteness of sampling using kick nets, resulting in low taxonomic overlap between water eDNA and kick net data from the same locations, which may also be beneficial depending on study objectives (Keck, Hürlemann, et al. 2022). Furthermore, one of the advantages of eDNA over bulk sample DNA is that eDNA can also be used for other taxa (e.g. invertebrates initially, followed by vertebrates, (Harper et al. 2021a), allowing for more efficient ecological surveys.

Overall, the eDNA-based methods detected a higher diversity than conventional sampling (up to double the number of genera in Uchida et al. 2020). However, both methods do not

always detect the same taxa (62% overlap between genera at regional scale, Mächler et al. 2019). Moreover, the target taxa are often a minor fraction in water eDNA datasets. Their proportion can increase by selecting the optimal primer pair (Leese et al. 2020, Brantschen et al. 2022; 2021) or by using multiple assays (Elbrecht and Leese 2017). On the other hand, the eDNA-based methods detected taxa such as small invertebrates and protists which were not detected by kick-net or other types of conventional sampling. eDNA methods result in a more holistic overview of freshwater biodiversity and more accurate assessment of ecological status at different trophic levels than conventional methods.

Despite the differences in community composition, both eDNA and conventional methods generated similar scores for freshwater biotic quality indices (Fernández et al. 2019; Seymour et al. 2020; Vivien et al. 2020). Some authors used machine learning to predict ecological status of rivers from eDNA data, when the number of traditional indicator taxa was low (Brantschen et al. 2021). A good congruence between metabarcoding data and morphology-based indices was also found for aquatic oligochaetes that are often used as bioindicators of sediment quality in rivers and lakes (Vivien, Lejzerowicz, and Pawlowski 2016). However, in this case the high throughput sequencing of individuals (DNA barcoding) was considered a more reliable method to infer the biotic indices because it provided abundance data (Vivien et al. 2020).

Few studies used qPCR rather than metabarcoding to detect invasive non-native (INNS) invertebrates (Dougherty et al. 2016; Blackman et al. 2020; Rice, Larson, and Taylor 2018). eDNA-based surveys using qPCR often outperform conventional kick-net sampling in detecting INNS, especially when the target species is present at low density. In contrast, eDNA-based surveys using metabarcoding sometimes detect fewer INNS than conventional sampling (Blackman et al. 2021). However, both approaches are complementary, as water eDNA analysis can detect species that are overlooked by traditional sampling.

4.3.3 Phytobenthos (Diatoms)

A total of nine publications were reviewed that investigated benthic diatoms as bioindicators of water quality in rivers and streams. Some of the papers focused on comparing diatom community composition in molecular and morphological datasets, others compare diatom indices derived from biofilm eDNA and microscopic data.

Several studies found discrepancies in taxonomic composition and abundance of diatom species inventories generated by microscopy and metabarcoding (Nistal-Garcia et al. 2020). These discrepancies have been explained by the gaps in reference databases, primer bias or issues in the bioinformatic pipelines (Bailet et al. 2019). The presence of cryptic diversity in many diatoms also complicates the taxonomic assignment of sequences. These discrepancies result in a mismatch between species found using morphological and eDNA-based methods. Species misidentifications may in turn influence the disparity between sequence-derived relative abundance and actual abundance, which becomes a source of the differences observed in ecological assessments (Zimmermann et al. 2014; Visco et al. 2015).

Solutions have been proposed to overcome these discrepancies to enable the use of diatom metabarcoding data in assessing water quality. Molecular indices were inferred from diatom data and their values were compared to morphological indices obtained from the same samples and used as a benchmark. The best results were obtained using taxonomy free approaches to predict molecular index or to assign indicator values to OTUs (Apothéloz-Perret-Gentil et al. 2021). A cell biovolume correction factor was proposed to overcome quantification biases (Vasselon et al. 2017). Its application generally improved the correlation between the two methods, suggesting that relative abundance estimates derived from sequence reads are feasible (Borrego-Ramos et al. 2021).

All reviewed studies agree that diatom metabarcoding has the potential to replace conventional microscopy-based analyses. However, given the important differences of diatom molecular and morphological data, this cannot be done by a simple replacement of specimens by sequences. Further research is needed to develop a novel metric derived directly from diatom metabarcoding data (M. Kelly et al. 2020).

4.4 Terrestrial

4.4.1 Vertebrates

A total of 33 publications were reviewed comparing conventional and eDNA-based approaches for detecting terrestrial and semi-aquatic vertebrates. Most of these papers used eDNA captured via water filtration followed by the detection of target species through qPCR. Soil eDNA, airborne DNA and invertebrate-derived DNA (iDNA) were also used to detect vertebrate species, although only a few comparative studies were available.

Amphibians and reptiles are common targets of eDNA-based surveys. For amphibians, the success rate for detection compared to conventional surveys can be very high (84% for newts, Rees et al. 2014). However, the number of taxa identified varies between the methods, with a large number of unique observations for both eDNA and conventional methods (Svenningsen, Pertoldi, and Bruhn 2022; Bálint et al. 2018). For some amphibians, the visual and audio techniques outperform metabarcoding, especially as individuals metamorphosed (Moss et al. 2022). Therefore, a combination of eDNA-based and conventional methods is recommended to correct for the detection biases inherent in surveys, especially when taking different life history traits into account.

Regarding reptiles, the performance of eDNA-based methods varies between studies (Lacoursière-Roussel et al. 2016; Davy, Kidd, and Wilson 2015; Piaggio et al. 2014). Conventional trapping outperformed qPCR assays for semi-aquatic snakes (Rose et al. 2019), but eDNA performed better for the detection of invasive pythons (Hunter et al. 2015). Thus, eDNA-based methods are useful for reptile surveys as part of general scoping, but not as a stand-alone method. The majority of studies targeted aquatic reptiles using eDNA from water samples, with less success in detecting reptiles from water eDNA than mammals (Nordstrom et al. 2022). This is widely considered to be due to the hard outer layer of reptile skin resulting in lower rates of DNA shedding (Adams et al. 2019). Studying terrestrial reptiles using soil eDNA or cover boards has not been extensively explored, but shows promise (Nordstrom et al. 2022; Kyle et al. 2022; Ryan et al. 2022; 2020). This has been shown to generate data for reptiles absent in camera traps, thus providing a complementary dataset (Ryan et al. 2022).

The effectiveness of eDNA surveys was also tested for mammal and bird monitoring. Mammalian eDNA was usually obtained from water or soil samples, but an alternative source was invertebrate-derived (iDNA) (Abrams et al. 2019, Gogarten et al. 2019) from insects and other invertebrates, such as mosquitos, flies, leeches or ticks, that have consumed or have attached mammalian DNA (see review by Carvalho et al. 2021). Bird eDNA is most commonly detected with broader vertebrate assays from water samples (Polanco Fernández, Mutis Martínezguerra, et al. 2021) even though there are bird-specific primers (Ushio et al. 2017). Birds have also been detected from eDNA collected from soils (Ryan et al. 2020), air (M. D. Johnson et al. 2023) and flowers (Newton et al. 2023).

Substrate plays an important role in eDNA surveys of terrestrial fauna. Sampling soil for vertebrate eDNA is limited in part by the unknown degradation timeframes of eDNA in soils (Leempoel, Hebert, and Hadly 2020) and has a lower detection probability than sampling water (Sales et al. 2020). This difference may be caused by the water mixing and spreading eDNA over a wider area than eDNA deposited in soils, although targeted sampling of soil around shelter locations may improve detections of particular species (Ryan et al. 2020).

This targeted sampling approach may be particularly viable when considering species distributions in arid areas (Egeter et al. 2018) or for species with a limited use of waterbodies (Harper et al. 2019).

In general, eDNA-based mammal surveys recover a similar number of taxa to conventional methods but, the composition of assemblages detected by eDNA differed from camera traps or mist nets. Camera traps generally detected larger taxa, while eDNA data was more effective in detecting smaller species (Mena et al. 2021; Harper et al. 2019; Sales et al. 2020; Lyet et al. 2021). The same trend was observed in the case of iDNA surveys, which preferentially detected large-bodied species compared to camera traps (Gogarten et al. 2020; Abrams et al. 2019).

In some environments, such as tropical forests, the reference database was a limiting factor for eDNA-based vertebrate surveys (Mena et al. 2021). The tropics are characterized by a high species diversity but there is relatively low financial support for DNA barcoding projects to increase the quality of reference databases. As reference databases expand to match those in temperate regions, so too will the capacity of eDNA-based metabarcoding for species level identification. A combination of eDNA-based and conventional methods is currently recommended in the tropics. The complementarity of both approaches for vertebrate monitoring in temperate regions is also recommended (Svenningsen, Pertoldi, and Bruhn 2022).

4.4.2 Terrestrial Invertebrates

A total of 12 publications were reviewed comparing eDNA-based methods to conventional methods in soil and above-ground invertebrate communities. The papers focused mainly on soil arthropods or nematodes, comparing their diversity in soil eDNA and bulk samples split for metabarcoding and morphological analysis.

The eDNA-based methods generally performed as well as or better than conventional methods when comparing terrestrial invertebrate communities. Most studies showed that metabarcoding detected a higher diversity of invertebrates than trapping or other conventional methods. However, the efficacy of soil eDNA metabarcoding depends on the taxonomic groups targeted in the study. eDNA-based methods outperformed morphological identification when targeting nematodes in terrestrial habitats (Kitagami and Matsuda 2022). However, Treonis et al. (2018) claim that metabarcoding can over- or underestimate the prevalence of some nematode families (Treonis et al. 2018). Morphology and metabarcoding of soil and bulk samples yielded well-correlated estimates of diversity and community composition of arthropods within soil (Oliverio et al. 2018). However, specific terrestrial invertebrate species considered as focal ecosystem service providers in orchards had very few taxa identified to a species level through the metabarcoding of soil eDNA (Todd et al. 2020). Similarly, metabarcoding revealed more species of termites in soil eDNA, but less species of ants and springtails compared to the morphological data of tropical soil fauna (Basset et al. 2022). Bulk sample metabarcoding can reproduce ecological patterns of morphologically identified beetle community from the same pitfall and malaise traps, but soil eDNA detected different beetle community and greater diversity of invertebrates than conventional methods (Watts et al. 2019). Due to the variability in results across taxa, multiple sampling methods are recommended for terrestrial insect biodiversity monitoring when carrying out DNA-based methods, as each has merits for particular ecological niches (Chua et al. 2023; M. Li et al. 2023).

4.5 Summary of Method Comparison

4.5.1 Fieldwork (Ease of Use, Sampling Effort, Cost, Safety)

Compared to the conventional methods, eDNA samples can be very simple to collect. This is particularly true in the case of water eDNA samples used to monitor fish and other aquatic (and semi-aquatic) vertebrates as well as aquatic invertebrates and microbiota. The main advantage of water eDNA sampling is that it can be done with far less expertise and effort and hence at much lower cost. Comparatively, the conventional methods take longer to obtain samples and often require trained ecologists or taxonomists to be present.

Water eDNA sampling is non-destructive and usually less hazardous, compared to trawling, netting or electrofishing. It can be more effective compared to visual observation and aerial surveys of marine vertebrates, especially when animals dive, or compared to PAMs (passive acoustic monitors), which detect mammals only when they generate noise. By contrast, for taxa shedding sufficient eDNA (e.g. mammals), they can be detected throughout the water column regardless of their activity (Booth, Sinclair, and Harwood 2020), provided the sampling strategy accounts for the dispersion and degradation of target eDNA.

There are some other practical advantages of using water eDNA applications to monitor industry activities in marine systems. For example, the towed equipment used to collect conventional samples is often impractical to use near marine infrastructure such as oil rigs, jetties, cables, pipelines, manifolds, or offshore windfarms due to the potential loss of gear, safety considerations and exclusion zones. These methods are therefore mainly applicable during the baselining phase, prior to construction of such infrastructure. If a follow-up survey is required once the facilities are in place as part of ongoing monitoring, conventional methods introduce additional health and safety exposure, hazards and overall execution risk. For continued monitoring of the impact of an offshore development, conventional imagery and eDNA methods present the best combination of options. In addition, eDNA will often detect elusive taxa that are living in or on marine structures, infrastructure, or artificial reefs, and which can be difficult or impossible to detect using imagery.

The water eDNA can be used to supplement other activities (ROV collecting data or visual inspection). To decrease vessel time and costs, the industry is using ROVs and AUVs for collecting a range of data, including visual inspection of the sub-sea infrastructure. Although equipment hire or purchase still represents a significant cost, ROVs and AUVs can also be used to collect eDNA samples (Closek et al. 2019; Scholin et al. 2017), imagery data (Alexander et al. 2022), and ichthyoplankton data (Pierella Karlusich, Lombard, et al. 2022). Furthermore, multiple eDNA samples can be collected over a temporal range for longer term monitoring using an automated sampler, compared to sampling during a single vessel deployment. For longer term surveys, autonomously sampled eDNA can be used (Searcy et al. 2022), which uses equipment similar to a fish trap and does not require regular field visits.

The sediment eDNA and bulk DNA sampling methods are broadly the same as conventional methods. However, a smaller volume of sediment is required for eDNA-based analyses, which makes it more suitable for sampling collection by AUVs and ROVs compared to conventional macrofaunal analysis. Also, less time is required for on deck processing (taking small cores or spooning subsamples from sediment grabs) saving considerable vessel time (Laroche et al. 2020a). The preservation of sediment and bulk eDNA samples has a significantly lower risk to safety, with minimal hazards compared to a formalin solution typically used for conventional sample preservation.

Regarding terrestrial sampling, the time and resources needed for a conventional field survey will vary significantly according to the size, accessibility and features of the site, as well as depending on the taxonomic targets. Camera surveys and some vertebrate trapping methods require the labour and resources for mobilising a large amount of equipment,

resulting in significant upfront and analysis costs (Mena et al. 2021). Sampling eDNA for terrestrial fauna typically takes less than half an hour per sample, not accounting for travel to the sampling location. Furthermore, this does not require a taxonomic expert for sampling in the field, but does require trained personnel for the subsequent analysis (Deiner et al. 2017). These advantages are less evident in the case of bulk DNA samples that are collected using the same methods as in conventional monitoring, whereby invertebrates are trapped, stored in buffer and then analysed via metabarcoding. Invertebrate communities can also be directly detected via soil eDNA sampling. An advantage of using soil samples is being able to survey for multiple taxonomic groups such as bacteria, fungi, and invertebrates simultaneously.

4.5.2 eDNA Sample Analysis (Processing Effort, Data Obtained, Reference Sequence Library, Turnaround Time)

The turnaround time is the main difference between conventional and eDNA analysis. Conventional methods can generate data instantly and in situ, for example during visual observations of marine megafauna or morphological identification of fish. Rapid identification can be critical when trying to adaptively manage industrial activities in the presence of a protected species, or when providing an early warning of the presence of INNS (Whitaker, Brower, and Janosik 2021). Although eDNA-based detections are becoming more rapid (Besson et al. 2022), they cannot generate immediate species observation data, hence conventional methods may be more appropriate where immediate detections are required.

On the other hand, eDNA-based methods can be automated and high-throughput, making them considerably less labour intensive and requiring less specialised taxonomist expertise compared to morphological analysis. This means that biological data can often be generated more rapidly and at a lower cost, especially when large number of samples is to be analysed or when many species need to be identified.

eDNA-based methods have been demonstrated to be comparatively cheaper than conventional methods with regards to labour and laboratory costs, while also requiring fewer in-field resources for the information gained. However, in some cases the eDNA analysis costs can be similar or even higher. For example, the costs of eDNA-based analysis per sample, can double the cost of camera trapping or small mammal traps due to the high unit costs. Yet, when comparing cost per species detected, eDNA-based methods have been shown to be a third of the price of camera trapping and just above half the cost of small mammal traps (Mena et al. 2021).

The ability to use multiplexing in the analysis of eDNA samples means that efficiencies and cost-savings are gained as the number of samples increases, whereas the processing of imagery or collected and sorted specimens increases proportionately to the total number of samples taken. Moreover, different taxa can be detected from the same eDNA samples, which reduces cost of sampling because DNA extraction is only done once per sample regardless of how many separate assays are run. Although reporting results from eDNA-based surveys may take longer, compared to some conventional surveys, they are more likely to include species of particular interest (e.g., INNS). The eDNA can also be used to examine ecosystem health more holistically and thereby may detect changes earlier than conventional methods, allowing for more timely management of industry activities.

4.5.3 Output Data

eDNA sampling consistently detects more taxa than conventional sampling, albeit with fewer identifications to species level, especially when using metabarcoding in marine and/or tropical environments. This is mainly due to the incompleteness of reference databases. The need to complete the reference database was highlighted as a priority by most of the reviewed papers.

The gaps in reference databases may also explain some of the disparity in species detected between eDNA and conventional methods. Taxonomic composition of metabarcoding data is broadly similar to morphology-based data, but their match largely depends on targeted taxa. The congruence is generally low for aquatic macroinvertebrates that are present in water or sediment eDNA samples only as DNA traces. The analysis of bulk samples provides more consistent results allowing detection of most species present in the sample. The taxonomic overlap of data from eDNA-based and conventional sampling can also increase with better primers.

On the other hand, an advantage of conventional methods is that abundance is usually included in the outputs as counts of individuals, which is more difficult to obtain with eDNA data. If population biological data such as size and abundance are required, conventional methods should be used. Although, advances are being made to overcome some of these challenges, for example, both qPCR and metabarcoding results frequently correlate with abundance and/or biomass for fish, this relationship is highly variable and direct inference is not robust (Luo et al. 2022). A cautious approach and regular calibration of eDNA-based methods against conventional methods is still needed when using eDNA for estimating abundance or biomass.

As remote and automated biological surveys become more prevalent, a combination of imagery (benthic and aerial), acoustic and eDNA data are likely to dominate environmental assessments. Broad species detections and community analyses from eDNA-based approaches can be complemented by the quantitative measurements generated by imagery data such as underwater video.

4.6 Summary of Advantages of eDNA-Based Methods vs Conventional Methods

As described in section 4.5, eDNA-based and conventional methods have each their strengths and limitations. The latter have been already listed in sections 2.2 and 3.3. The advantages of each are summarized in Table 4.1 with reference to the fieldwork, sample analysis and output data.

Table 4.1 – Advantages of eDNA-based biomonitoring and conventional methods at different stages

	Advantages eDNA	Advantages conventional
Field work	<ul style="list-style-type: none"> • Samples easy to collect, require less time and limited expertise • Sampling more effective - less dependent on species activity and habitat • Samples can be obtained in parallel to other monitoring activities • No need to use towed equipment near industrial installations • Sampling non-invasive • Samples preservation using non-hazardous solutions 	<ul style="list-style-type: none"> • Visual observations, netting or trapping provide instant, real-time data • Camera traps and other trapping devices have comparable low unit costs • Sampling protocols are standardised for different habitat and substrate types
Sample analysis	<ul style="list-style-type: none"> • High-throughput automated sample processing • Reduced time and costs when many samples processed simultaneously • No need for taxonomic expertise • Same sample types can be used for surveying multiple taxonomic groups 	<ul style="list-style-type: none"> • Turnaround time of sample processing and storage can be rapid • Sample processing requires a relatively simple equipment

Output data	<ul style="list-style-type: none"> • More taxa consistently detected • More sensitive to detect rare species • Broader taxon range data available • Possible identification of inconspicuous and cryptic taxa • Species identification less subjective, depending solely on taxonomic coverage of reference database • Retroactive analyses possible 	<ul style="list-style-type: none"> • Species Identification possible if taxonomic expertise available • Abundance data obtained through counting specimens • Biomass, age and health state can be reported • Biotic indices well established
--------------------	--	--

4.7 Recommendations

Based on the reviewed papers, the status of eDNA-based methods compared to conventional methods was established. The criteria taken into consideration were the complexity, costs and health and safety risks of field sampling, the turnaround time and costs of sample processing and comprehensiveness of output data. For each habitat and target taxa, the eDNA-based methods were evaluated as:

- Stand-alone tools
- Preferentially used in combination with conventional methods
- Technically feasible but requiring further testing and validation

When appropriate, the single species detection and community analysis were evaluated separately. We assume that the appropriate markers for species detection are available.

Table 4.2 – eDNA method status for different taxa and data analyses

Habitat	Taxon	eDNA analysis	eDNA method status
Marine	Fish	Community analysis	Stand-alone
		Species detection	Stand-alone
	Other vertebrates	Community analysis	In combination
		Species detection	In combination
	Soft bottom benthos	Community analysis	Further validation
	Hard bottom benthos	Community analysis	In combination
	Zooplankton/Phytoplankton	Community analysis	In combination
		Species detection	Stand-alone
Freshwater	Fish	Community analysis	In combination
		Species detection	Stand-alone
	Aquatic invertebrates	Community analysis	Further validation
		Species detection	Stand-alone
	Phytoplankton (diatoms)	Community analysis	Further validation
Terrestrial	Vertebrates	Community analysis	In combination
		Species detection	In combination
	Soil invertebrates	Community analysis	Further validation
	Above-ground invertebrates	Community analysis	In combination

5 Application of eDNA-based Surveys in the Energy Industry

5.1 Review of Existing Literature

Numerous studies explored the potential of eDNA-based technology to assess the environmental impact of energy project activity. The topics of these studies include:

- Monitoring the impacts of exploratory drilling and discharges from offshore platforms
- Monitoring the impacts associated with hydraulic fracturing
- Monitoring the impacts of oil spills
- Assessing long-term effects of decommissioning

Among 29 papers reviewed, ten papers used sediment metabarcoding to assess the environmental impact of offshore platforms. The focus of these studies was to find new bioindicators among microbial and meiofaunal communities that could potentially replace macrofauna currently used in routine benthic monitoring (Lanzén et al. 2016; Mauffrey et al. 2020). Laroche et al. (2017) showed that bacterial communities exhibited the strongest response to drilling and extraction impacts, followed by foraminifera, while macroinvertebrates (identified using morphology) were the least responsive. The microbial community was also a focus of the study by Krolicka et al. (2020), who found that the composition and abundance of bacterial taxa correlate with the level of petroleum-related compounds such as barium and PAH. Correlation with environmental parameters (barium, THC and heavy metals) was demonstrated in most of benthic eukaryotic communities present in the vicinity of offshore oil and gas platforms (Laroche et al. 2016; Frontalini et al. 2020). Both water and sediment metabarcoding show that the impact is limited to the area less than 200m from the platform (Cordier 2019). Eukaryotic metabarcoding data were also used to calculate biotic indices, which were shown to be consistent with those inferred from benthic macrofauna (Mauffrey et al. 2020).

Environmental genomic methods were also used extensively to assess the effects of oil spills and oil pollution on marine coastal habitats. Metabarcoding, metagenomic and metatranscriptomic studies analysed the impact of Deepwater Horizon spill on bacterial and eukaryotic communities (Bik et al. 2012; Mason et al. 2012). Long-term ecological effects of residual oil on benthic communities were analysed in a contaminated area on the coast of South Korea (Xie et al. 2018), while Oladi et al. (2022) analysed the impact of frequent crude oil spills on coral reef ecosystem in the Persian Gulf. The mesocosm experiments conducted by Krolicka et al. (2017; 2019) allowed for the identification of key microbial indicators of crude oil pollution, while Knapik et al. (2020) used metatranscriptomics to identify functional genes in bacterial communities that could be used as target for oil-related genosensing.

Compared to many studies applying eDNA to the marine environment, much less work has been done on monitoring energy project activity in freshwater and terrestrial environments. Few studies were conducted on the impact of hydraulic fracturing on microbial communities living in the streams, however, they were limited to northwestern Pennsylvania, and only investigated bacterial communities (Ulrich et al. 2018). Bacterial metabarcoding was also used to test different soil covers used for the decommissioning of oil sands mining sites (F. Stefani et al. 2018). On the other hand, freshwater zooplankton metabarcoding was used to assess the efficiency of oil spill remediation activities in a boreal lake (Ankley et al. 2021). Additionally, water eDNA has been used to evaluate the impact of a gasoline spill on populations of giant salamander in Pennsylvania (Perelman et al. 2021).

Table 5.1 – eDNA studies of energy project activity and related impacts

O&G activity	Target taxa	eDNA method	Project objective	Reference
Exploratory drilling	Bacteria	Sediment metabarcoding	Microbial bioindicators of drilling waste discharge	Nguyen et al. (2018)
Exploratory drilling and gas production field	Bacteria	Sediment metabarcoding	Bacterial community metabolic response	Laroche et al. (2018)
Oil platforms (New Zealand)	Bacteria and eukaryotes	Sediment eDNA and eRNA	Benthic monitoring of offshore drilling sites	Laroche et al. (2017)
Oil platforms (New Zealand)	Benthic foraminifera	Sediment metabarcoding	Macrofauna and foraminifera response	Laroche et al. (2016)
Oil platforms (North Sea)	Eukaryotes	Phylogenetic microarray	Benthic monitoring of oil-drilling sites	Lekang et al. (2020)
Oil platforms (North Sea)	Eukaryotes	Sediment metabarcoding	Benthic monitoring of oil-drilling sites	Lanzen et al. (2016)
Oil platforms (North Sea)	Bacteria and eukaryotes	qPCR and metabarcoding	Identification of microfaunal benthic indicators	Krolicka et al. (2020)
Oil platforms (North Sea)	Eukaryotes (3 markers)	Sediment metabarcoding	Biotic indices inference	Mauffrey et al. (2020)
Oil platforms (Persian Gulf)	Elasmobranch	qPCR, water metabarcoding	Whale shark aggregation at an oil field	Thomsen et al. (2016)
Gas platforms (Adriatic)	Eukaryotes (5 markers)	Water and sediment metabarcoding	Response of planktonic and benthic communities to platforms activity	Cordier et al. (2019)
Gas platforms (Adriatic)	Benthic foraminifera	Sediment metabarcoding	Comparing morphology and metabarcoding data	Frontalini et al. (2020)
Oil spill (Gulf of Mexico)	Eukaryotes	Sediment metabarcoding	Deepwater Horizon oil spill impact on coastal habitats	Bik et al. (2012)
Oil spill (Gulf of Mexico)	Bacteria	Metabarcoding, Metatranscriptomics	Oil spill impact on beach microbial communities	Lamendella et al. (2014)
Oil spill (Gulf of Mexico)	Bacteria	Metagenomics, Metatranscriptomics	Response of bacterial communities to aliphatic hydrocarbons	Mason et al. (2012)
Oil spill (Gulf of Mexico)	Nematodes	Bulk DNA metabarcoding	Oil spill impacts on intertidal meiofauna	Brannock et al. (2014; 2017)
Oil spill (Persian Gulf)	Bacteria and eukaryotes	Sediment metabarcoding	Oil spill Impact on coral reef sites	Oladi et al. (2022)
Oil spill (South Korea)	Bacteria and eukaryotes	Sediment metabarcoding	Microbial and metazoan response to oil spill	Xie et al. (2018)

Gasoline spill (river)	Amphibians	Water eDNA	Impact on endangered giant salamander	Perelman et al. (2021)
Oil spill remediation	Freshwater zooplankton	DNA and RNA metabarcoding (bulk samples)	Natural vs active shoreline cleaning in freshwater ecosystems	Ankley et al. (2021)
Oil spill remediation	Freshwater zooplankton	DNA and RNA metabarcoding (bulk samples)	Experimental testing of zooplankton response to environmental stressor	Ankley et al. (2022)
Oil pollution	Bacteria	qPCR and metabarcoding	Identification of microbial key indicators	Krolicka et al. (2019)
Oil pollution	Bacteria	Water metabarcoding	Mesocosm experiment	Krolicka et al. (2017)
Oil pollution	Bacteria	Meta-transcriptomics	Development of genosensors	Knapik et al. (2020)
Hydraulic fracturing	Bacteria	Water, biofilm and sediment metabarcoding	Response of stream microbial communities to environmental impacts	Trexler et al. (2014)
Hydraulic fracturing	Bacteria	Biofilm metabarcoding	Chemical pollution impact on streams microbiota	Johnson et al. (2017)
Hydraulic fracturing	Bacteria	Water and sediment metabarcoding	Response of stream microbial communities to environmental impacts	Ulrich et al. (2018)
Decommission oil sands	Bacteria and fungi	Soil metabarcoding	Impact of soil covers on oil sands mining sites	Stefani et al. (2018)
Decommission oil platforms	Fish and elasmobranch	Water metabarcoding	eDNA vs ROV stereo-video observations	Alexander et al. (2022)
Biocorrosion control	Bacteria	Water metabarcoding	Monitoring microbiota in produced water	Dutra et al. (2023)

5.2 Conventional vs eDNA-Based Monitoring Using Offshore Platforms as a Model

Here, we present the current conventional methods used for biomonitoring of offshore platforms and discuss the potential of eDNA analyses to replace and/or complement these methods as the primary model for consideration. Our selection of conventional methods is based on the *Guidelines for environmental monitoring of petroleum activities on the Norwegian continental shelf* (Norwegian Environmental Agency 2020), to which we added acoustic and visual observation of marine mammals from the UK JNCC Guidelines during geophysical operations (JNCC 2017).

Currently, the only biological component included in water column monitoring of offshore platforms are ecotoxicological analyses of contaminants present in tissues of wild caught fish and caged blue mussel. Monitoring the communities of fish and other pelagic species is not considered in the current regulations. However, the Guidelines will consider using eDNA methods as part of water column monitoring in the near future. An example of how these tools could be applied is in the combined use of eDNA with acoustic and visual surveys to detect cetaceans during baseline surveys. In addition, water eDNA metabarcoding could also be used to periodically survey the impact of energy project activity on pelagic biota.

With regards to seabed monitoring, conventional surveys include both hard- and soft-bottom macro- and megafaunal diversity. Softbottom meiofauna (e.g., foraminifera, nematodes) are also mentioned, though they are considered as an additional activity. Visual surveys using ROV video footage are proposed for monitoring hardbottom fauna (e.g., corals, sponge and sea-pen communities), whereas softbottom macrofauna are analysed based on sediment grab samples. In both cases, the morphological species identification could potentially be replaced by eDNA, either by analysis of DNA extracted from bulk samples or eDNA metabarcoding of water or sediment samples. Both methods can be used to provide species list and taxonomic composition of benthic communities. As described above, sediment metabarcoding could also be used to assess ecological status of benthic communities, with a focus on microbial or meiofaunal bioindicators.

Table 5.2 – Conventional monitoring methods of energy project activity in marine environment and corresponding eDNA-based solutions.

Conventional monitoring (operation phase)	Potential eDNA solution
Water column monitoring	
Marine mammals (cetaceans): acoustic and visual observation	Water eDNA metabarcoding or qPCR
Wild caught fish tissue ecotoxicological analysis	Not applicable
Blue mussel contaminants monitoring	Not applicable
Benthic monitoring	
Hardbottom benthic megafauna: visual survey, mapping of vulnerable species habitats	Bulk DNA or water eDNA metabarcoding
Softbottom benthic macrofauna: morphology-based identification, counting, diversity indices	Bulk DNA or sediment eDNA metabarcoding
Softbottom benthic meiofauna: morphology-based identification, diversity indices	Sediment eDNA metabarcoding

5.3 Prospective eDNA Applications to Monitor Energy Project Activities

Here, we propose some eDNA applications that could be relatively easy to implement at different phases of energy project lifecycle. We focus on marine environments, where most of the referenced studies relating to energy project activities were conducted, but these applications could be easily transferred to the freshwater and terrestrial habitats. The current development status of each eDNA method is provided based on the reviewed papers (section 5.1).

Table 5.3 – Prospective eDNA methods to assess the ecological impacts of energy project activities in marine environments

eDNA method	Target taxa	Aims and Expected results	Current status
Baseline surveys			
Water eDNA metabarcoding/qPCR	Marine mammals (cetaceans)	Avoid disturbance and inference with migratory routes for selected sites	Ready to use
Water eDNA metabarcoding	Fish/pelagic macrofauna	Detect sensitive species / avoid fishing spots	Ready to use
Bulk DNA metabarcoding	Zooplankton	Surface water plankton taxonomic composition	Development phase
	Benthic macrofauna	Benthic macrofauna taxonomic composition	Development phase
Sediment eDNA metabarcoding	Benthic meiofauna	Meiofauna taxonomic composition	Development phase
	Benthic microbial community	Microbial taxonomic composition	Ready to use
Exploratory and production drilling			
Water eDNA metabarcoding	Bottom fish and other fauna	Assess bottom water biological quality	Development phase
Sediment eDNA metabarcoding	Benthic meiofauna and microbial community	Assess sediment biological quality	Development phase
Oil spills and remediation			
Water/sediment eDNA metabarcoding	Microbial community	Assess the impact of contaminants	Ready to use
Water eDNA metabarcoding/qPCR	Fish/pelagic macrofauna	Detect sensitive species	Ready to use
Decommissioning			
Bulk DNA metabarcoding	Hardbottom invertebrates	Assess epibenthic diversity	Development phase
Water eDNA metabarcoding/qPCR	Invasive non-native species (INNS)	Detection of INNS	Ready to use
Restoration			
Bulk DNA metabarcoding	Benthic macrofauna	Congruence with baseline studies	Development phase
Water eDNA metabarcoding/qPCR	Fish/pelagic macrofauna	Detect sensitive species, reintroduction of species	Ready to use
Paleogenomics – sediment ancient DNA	Macrophytes, macrofauna	Reconstruct reference conditions of impacted ecosystems	Development phase

5.3.1 Baseline Surveys

Baseline assessments provide a starting point from which to measure potential environmental change. A baseline often consists of a survey of an existing habitat or ecosystem, which may experience subsequent ecological change from energy project activities. These are static baselines that can be used to illustrate change over time. For determining if changes are specific to the location, a dynamic baseline which accounts for change over time is generally recommended. Although this is reliant upon information over a wider area, including reference stations, this design allows any change to be given a broader context.

Environmental DNA-based approaches are particularly useful during the baselining stage because they offer a rapid and relatively complete overview of what is living around prospective energy project activities. Compared to conventional methods, eDNA approaches do not require extensive knowledge of each taxonomic group but can provide an inventory of a wide range of taxa, depending on their representation in reference databases. This unique insight into global biodiversity can be obtained based on relatively few water and sediment samples, as the diversity of different taxonomic groups can be explored using the same eDNA samples, saving time and costs during sample collection and processing.

Environmental DNA can also be used to detect key species, such as protected or sensitive species, which is critical during baselining. It can inform actions such as the complete avoidance of particular areas, or the avoidance of activity during certain times of year, which is the first stage of the mitigation hierarchy. The monitoring of marine mammals and their migratory routes is particularly important with regards to the location of offshore energy activities and infrastructure. Migratory routes are currently monitored using acoustics, aerial imagery and visual observations but this could easily be complemented by eDNA-based approaches. Conducting a broad baselining study using these complementary methods would be less invasive and cover a range of temporal and spatial scales. Additionally, eDNA could also be used for the survey of particularly sensitive habitats, such as spawning areas or coral reefs. These eDNA approaches are well established for detecting pelagic species in the water column, but, they could also be applied to hard bottom fauna in combination with ROV video imagery.

There are already examples of using eDNA-based approaches for baseline surveys in marine industrial activities other than those associated with the energy project activities. For example, eDNA was used to explore deep-sea biodiversity in the area of prospective polymetallic nodule mining (Laroche et al. 2020a; 2020b). The studies show that benthic metazoan diversity is positively correlated with nodules density and sea bottom topography and that the pelagic eDNA does not impact surveys of benthic community at the deep-sea floor. This type of water and sediment eDNA metabarcoding analyses are technically ready to use and could be easily implemented also during the pre-production baselining of energy industry activities.

5.3.2 Operations Phase

As shown in section 5.1, several studies applied eDNA-based methods for biodiversity monitoring during the operational phase of energy projects. Most of them focus on the assessment of impacts associated with the activity of offshore platforms on benthic diversity. Traditionally, this is done based on benthic macrofauna sorted from sediment samples, morphologically identified and counted. The macrofauna data are used to infer biotic indices, such as AMBI (Borja, Franco, and Pérez 2000) or BQI (Benthic Quality Index, Rosenberg et al. 2004), that serve to assess the ecological status of a benthic community.

Inferring macrofauna-based indices from bulk or sediment eDNA metabarcoding is possible (Lejzerowicz et al. 2015) but not always reliable due to the issues of abundance, variation in

biomass and low representation of macrofauna in small sediment eDNA samples (Pawlowski et al. 2022; Wort et al. 2022). These limitations can be overcome by predicting biotic indices from bacterial or eukaryotic eDNA metabarcoding data using machine learning approach (Cordier et al. 2018). Yet, to make this approach successful an extensive training dataset are needed. Alternatively, the macrofauna can be replaced by meiofauna or microbial indicators and indices based on sensitivity to drilling-related contaminants (Laroche et al. 2016; T. T. Nguyen, Cochrane, and Landfald 2018; Krolicka et al. 2019)

5.3.3 Oil Spills and Remediation

Monitoring oil spills is of key importance for the sustainability of the marine environment. When an unplanned event has occurred, such as a chemical/waste leakage or an oil spill, an environmental survey may be required to comply with a regulatory requested environmental impact assessment to establish a post-impact baseline (Hinz et al. 2022). Different sensor technologies are available to detect and monitor oil spills. Remote sensing data can be used to characterize oil spill pollution types (Yang et al. 2023). The environmental impact of oil spills can also be assessed based on microbial communities, particularly bacteria and fungi, as these organisms have rapid responses to environmental changes. The protocols for eDNA metabarcoding of bacterial and eukaryotic microbiota are well-established, and the methods are ready to be applied.

Monitoring microbial communities also plays an important role during the remediation process. Selected bacterial groups can be used as indicators of oil pollution (Krolicka et al. 2019) and some of bacterial genes are considered as promising biosensors (Krolicka, Gomiero, and Baussant 2020). Furthermore, the eDNA metabarcoding and metagenomic data could also assess ongoing remediation by targeting microorganisms that support biodegradation and assimilation of hydrocarbons as well as screening for hydrocarbon-degrading genes. Bacterial community changes can indicate the progression of oil spill degradation.

Alternatively, eDNA-based surveys can be used to rapidly obtain samples for a snapshot of the ecosystem condition, potentially even between the spill and the impact. These samples can be archived in a biobank to provide a reference for future remediation work, or simply be used to monitor the recovery of the impacted site (Yergeau et al. 2015).

5.3.4 Decommissioning

At the end of production phase, decommissioning is the first step towards returning the site to its natural pre-production conditions. Depending on the regulations, the installations can be entirely removed or part of them can be left in place. If the installation is not completely removed, it will continue to act as an artificial substrate. Such substrate can be rapidly colonized by various epibenthic species and become a hotspot of diversity, which impacts the whole benthic community in platform vicinity. The species living on such hard substrate belong to various groups of sessile invertebrates, whose identification using visual observation and morphology-based taxonomy is often problematic.

Environmental DNA metabarcoding appears as the best solution to survey a huge diversity of sessile organisms living on hard substrates (Obst et al. 2020). Several studies demonstrate its efficiency, as in the case of Autonomous Reef Monitoring Structures (ARMS) (Pearman et al. 2020; Ip et al. 2023). The significant increase of benthic diversity was also demonstrated using morphological and metabarcoding analyses in the case of a partly decommissioned gas platform in the North Sea (Klunder et al. 2018). The impact of decommissioned platforms on marine diversity in the Gulf of Thailand was analysed through eDNA metabarcoding of water to target fish and elasmobranchs (Alexander et al. 2022), as well as to explore broad range of eukaryotic phyla using water, bio-foul and sediment eDNA samples (Alexander, Marnane, McDonald, et al. 2023). Finally, eDNA metabarcoding has

also been used to detect marine invasive non-native species (INNS) (Zaiko et al. 2020). The partly decommissioned platforms could act as steppingstones for INNS dispersal, therefore it is of great ecological importance to include their detection in routine monitoring (Macreadie, Fowler, and Booth 2011).

5.3.5 Restoration

Ecological restoration is the practice of renewing or restoring degraded, damaged, or destroyed ecosystems and habitats via active human intervention. Monitoring diversity before, during and after restoration efforts is used to assess changes in these ecosystems. Conventional approaches for monitoring restoration include visual surveys, trapping/netting, tissue sampling, and sampling for chemical analysis, such as heavy metals and hydrocarbons.

Environmental DNA-based methods offer the potential to assess changes to the biological communities compared to the original baseline. One of the advantages is that the eDNA sampling method can remain constant throughout the changes to the habitat, whereas most forms of conventional sampling would have to be altered due to changes in the habitat type. For example, new artificial reef structures make net tows for plankton or fish prone to entanglement. Similarly, saltmarshes restored on bare substrate make deployment of underwater cameras for fish community analyses more problematic, whilst a water eDNA sample remains feasible. Furthermore, eDNA-based methods generally have higher detection rates for cryptic and elusive species than conventional methods, and the capability to sample a broader community. This makes eDNA-based methods very useful when tracking the reintroduction of rare species (Riaz et al. 2020; Rojahn, Gleeson, and Furlan 2018) or monitoring any changes to the local biodiversity and ecosystem condition following restoration (Armbruster et al. 2021; van der Heyde, Bunce, and Nevill 2022; Heyde et al. 2020). These benefits of eDNA-based monitoring can also be achieved at a lower effort, increasing overall project efficiency in restoration monitoring.

The use of eDNA-methods applied to restoration can be limited by the lack of genetic data for pre-construction baselining. Various technical and biological factors can also bias the comparison of baseline surveys recorded using conventional and eDNA methods. However, this can be overcome using eDNA data from closely situated, non-impacted reference sites. Alternatively, it is possible to recover reference conditions through the paleogenomic analysis of sedimentary archives. Recently, environmental paleogenomics was used to investigate the impact of industrialization, urbanization and agriculture development during Anthropocene (e.g. Barrenechea Angeles et al. 2023; Siano et al. 2021). Such eDNA studies conducted on decadal scale could be of great interest to recover reference conditions at decommissioned sites and other areas where industrial activity has stopped but the past biodiversity has not been well documented (N.-L. Nguyen et al. 2023).

6 Elements to Consider When Planning eDNA-Based Surveys

This section discusses elements that should be considered when planning to include eDNA in biomonitoring surveys, as a replacement or complement to conventional methods. As shown in the preceding sections, the potential of an eDNA-based approach for the biomonitoring of energy project activities is high. Yet, eDNA approaches and their results may not always meet the criteria and requirements of biomonitoring procedures. Hence, it is important to carefully assess the values of an eDNA approach compared to conventional methods before its practical application. For each element described below, we provide some guiding questions to be asked when selecting the best approach.

6.1 Suitability

The first element for consideration is the capacity of eDNA-based approaches to fulfil the purpose of a specific biomonitoring activity. In general, eDNA provides information about the presence/absence or relative abundance of organisms in the sampled environment. It allows the identification of the species, or group of species based on their DNA. Hence, it is a valuable tool for detecting target species, for example when monitoring the presence of marine mammals during baseline surveys or searching for invasive non-native species (INNS) during the operational phase. The use of eDNA as a replacement or complement to conventional methods is also possible if the goal of the survey is to analyse community response to potential impacts associated with energy project activities, (e.g. benthic monitoring around offshore platforms). It is possible to replace morphotaxonomy through eDNA-based community analysis which may be beneficial for ecosystem health assessment, however, this remains dependent upon a good understanding of DNA sources, biases and error levels as well as sample representativity, as demonstrated in Lanzén et al., (2021).

On the other hand, eDNA-based methods cannot usually be used to biologically characterize the populations to inform directly about the age (although see Zhao, van Bodegom, and Trimbos 2022), biomass, or metabolic activity of target taxa. Some of this information could be obtained indirectly by analysing the eRNA data but the interpretation of such data remains problematic (Cristescu 2019). Nor can an eDNA approach currently contribute substantially to ecotoxicological tests that are commonly used to assess the impact of pollutants associated with energy project activities. This also concerns the analysis of DNA adducts that can be used for the detection of genotoxic chemicals in the environment (Pampanin et al. 2017). In all these cases, the conventional methods are more suitable.

- Questions:
 - How suitable is the eDNA method to fulfil the objectives of the survey?
 - How well will the eDNA-based biodiversity assessment inform about the environmental changes?
 - What kind of biological data is required by the survey?

6.2 Feasibility

The second element is to ensure whether there are any technical challenges that could preclude the use and processing of eDNA samples. Compared to conventional methods, an eDNA-based approach is often viewed as technically more complex, and this could discourage its potential users. Sampling for eDNA can be faster and easier, but some precautions must be taken which are usually not required by conventional biomonitoring. The processing of eDNA samples requires access to laboratory infrastructure and a sequencing facility. Therefore, it is important to ensure that eDNA sampling can be conducted in optimal conditions and that molecular facilities are available without triggering issues around logistic or regulatory constraints when sending samples.

The feasibility also depends on how easy it is to detect the species or taxa of interest through eDNA. Gaps in reference databases may also impact the accurate taxonomic assignment of sequences and can also affect the ability to design species-specific assays (Weigand et al. 2019). Moreover, some taxa may be difficult to detect if they only shed small amounts of DNA into the environment. If this is the case, it might be easier to achieve monitoring goals using conventional methods. Alternatively, eDNA could be used as to complement visual observations or other monitoring tools.

- Questions:
 - Are the eDNA protocols well established and is the necessary equipment available?

- Are the results of eDNA surveys easy to interpret?
- Are the reference sequences of target species present in the database?

6.3 Reliability

The third element to be considered when selecting the method is how trustworthy the eDNA analysis is. Most conventional methods are based on visual observations and morphological identification. The efficiency and accuracy of field observers and laboratory staff depends on their personal skills and taxonomic knowledge, which is often insufficient to correctly identify specimens to the species level.

An eDNA-based approach, especially when based on traces of macro- and megafauna, provides indirect evidence of the presence/absence of a target species. There are some technical and biological factors (transportation, degradation, etc.) that can influence the results of eDNA analysis. It is important to ensure that the interpretation of eDNA results accounts for these potential biases. On the other hand, eDNA analysis is more reliable in terms of species identification, which is based on automatic assignment of the sequences to the reference database. Its efficiency depends on the completeness of the database and the taxonomic resolution of the DNA barcode. These factors must be taken into consideration when eDNA analyses are planned.

- Questions:
 - Do the eDNA results match those of conventional methods for target species?
 - Are the potential biases of eDNA analyses accounted for?
 - How good is the taxonomic resolution of the selected marker?

6.4 Acceptance

The validation and acceptance of eDNA methods is another factor to be carefully appraised. In general, conventional biomonitoring is based on well-known bioindicator taxa, whose ecology has been extensively studied. Most of the conventional methods have been standardised and are accepted by official regulations (e.g. WFD, MSFD, Water Act, etc).

The formal acceptance of eDNA-based methods for use in routine biomonitoring is still ongoing. Although many eDNA applications have been scientifically tested, their validation in rigorous conditions has rarely been accomplished. The analysis of 546 published single-species assays show that the majority of investigated assays are incomplete (Thalinger et al. 2021). Similarly, the eDNA-based assessments of ecological quality status often lack rigorous validation, such as the ring-tests conducted for benthic diatoms monitoring (Vasselon et al. 2021).

Nevertheless, although only formally accepted for limited applications, an eDNA approach is usually considered a valuable alternative by regulators and environmental protection agencies. When choosing between conventional and eDNA methods it might be important to check whether the results of eDNA analysis could be accepted even if the method is not formally recognized. This is often not a problem if the eDNA approach presents a significant improvement of conventional methods in terms of sensitivity and quality. Acceptance can be more difficult if the eDNA approach directly competes with the conventional methods, often due to the reluctance of those that use these methods (e.g. diatoms-based assessment of the ecological quality of rivers and streams). To overcome this resistance to change, more convincing demonstrations of the effectiveness of these eDNA applications may be needed.

- Questions:
 - Is the planned eDNA method scientifically supported?
 - Is the method accepted by regulators?

6.5 Economic advantages

An eDNA-based approach presents some important economic advantages compared to conventional methods. It is generally expected that eDNA-based approaches will reduce costs considerably. A comparative study of costs related to deep-sea biodiversity assessment indicate that eDNA metabarcoding is about one tenth the cost of conventional methods using morphological taxonomy (Le et al. 2022). It is anticipated that these costs will rapidly decrease as sequencing technologies continue to develop. However, it is important to keep in mind that the costs of eDNA analyses are reduced mainly due to multiplexing, which consists of processing large numbers of samples at the same time. A conventional approach may still be a financially comparable option for smaller projects, where fewer samples are analysed. A conventional approach could also be a solution in countries where the labour costs are low, and taxonomic expertise is available.

Environmental DNA approaches can also allow for important time savings. In the case of marine monitoring, the sampling can be done more rapidly saving ship time, reducing costs considerably. Sediment sampling for eDNA is by far faster than sieving and sorting benthic macrofauna. Similarly, collecting water samples for eDNA analysis takes less time than fishing or zooplankton sampling. The turnaround time to results is another aspect to consider. While it can take up to 3 months to identify and count all the members of a pelagic or benthic community through conventional methods, the analysis of eDNA from water or sediment samples can take less than 2 weeks. The faster potential turnaround for analysis may be useful if an unplanned event occurs or if a rapid certification is needed. On the other hand, the time factor may not be as important in the case of routine monitoring that occurs yearly or every second year, and the results of which have no direct impact on energy industry operations.

- Questions:
 - Does the use of eDNA approaches allow for a reduction in costs?
 - Does it allow for time savings?
 - Is time an important factor for the study objectives, operational considerations or decisions?

6.6 Added value

Despite financial considerations, it is also important to consider the added value of eDNA-based approaches compared to conventional methods. First, the sensitivity of eDNA assays should be considered. There is a general agreement that the eDNA method is more sensitive because it can detect traces that organisms left in the environment. This is particularly important when the species are rare and difficult to observe, as is often the case in the detection of endangered species or as an early warning of biological invasions.

Another important added value is the holistic nature of eDNA data. An eDNA-based approach can provide insights into global biodiversity, encompassing a wide range of taxa with equal or minimal effort, including the most inconspicuous components of biological communities that could never be explored using conventional methods. In principle, all taxa can be surveyed using eDNA, while morphology-based identification is generally limited to a few generally large-sized taxa. It is also important to highlight that the eDNA approach can generate large amounts of data that can be easily integrated into a global network of information. This can considerably improve biomonitoring by giving access to comparable data in temporal and spatial dimensions, also allowing for the potential retroactive analysis across the life cycle of the project. Further automation of eDNA methods will allow the monitoring of the status of the environment in real time and worldwide.

- Questions:

- Is the eDNA-based approach more sensitive than conventional methods regarding target taxa?
- Can the holistic assessment of biodiversity make monitoring more efficient?
- Can the retroactive analysis of eDNA data be useful for the project?

6.7 SWOT Analysis

In this section we propose a SWOT analysis comparing Strengths, Weaknesses, Opportunities and Threats of eDNA projects versus conventional biomonitoring methods. Table 6.7 summarizes the pros and cons that should be considered when selecting the survey method.

Table 6.1 – SWOT analysis of eDNA-based biomonitoring vs conventional methods

Strengths	Weaknesses
<ul style="list-style-type: none"> ○ Easier sampling logistics: short training time, reduced hands-on time, less specialized equipment ○ Non-invasive sampling ○ Automation: the samples are processed automatically following established standards ○ Speed: large number of samples can be processed in a short time ○ Low cost: processing of eDNA samples can often be less expensive ○ Objective results: personal taxonomic expertise is not required ○ High sensitivity: allows detection of rare species 	<ul style="list-style-type: none"> ○ Biological data: Some biological data such as population age, biomass, abundance, abnormal growth cannot be obtained from eDNA ○ Localization: it may not be possible to obtain a precise location for species due to eDNA movement within the environment ○ Gaps in barcoding database can impede species identification ○ Limited DNA shedding can make some species difficult to detect ○ Technical and biological biases are not always well understood, limiting quantitative interpretation of eDNA data ○ Sampling method that would be optimal for species detection is not always feasible (e.g.no water at a site)
Opportunities	Threats:
<ul style="list-style-type: none"> ○ Environmental regulations: allows to respond to constantly increasing demand for biomonitoring. ○ Global biodiversity changes: wide range of taxa can be identified ○ Innovative technology: the methods are constantly improving ○ Retroactive analysis: the eDNA samples can be stored and reanalysed with different markers if the need arises 	<ul style="list-style-type: none"> ○ Acceptance: new technology can take long to be validated and adopted by regulators ○ Regulations: biomonitoring requirements are highly diversified between countries

6.8 Decision Tree for Selecting eDNA-Based Methods for Biomonitoring

To conclude, we propose considering the following decision tree when planning a biomonitoring project and choosing between eDNA and any conventional methods.




Figure 5. Decision tree for choosing between eDNA-based or conventional methods for species identification and biodiversity assessments

7 Future Directions of eDNA Methodologies

7.1 General

Environmental DNA-based methods have the potential to make environmental surveying safer and easier, more efficient, and provide more actionable results. This section gives a brief overview of how developments in key areas may allow adoption of eDNA-based methods alone, or along with conventional methods.

7.2 Sampling Techniques, Equipment, and Surveying

7.2.1 Automated sampling

The deployment of automated eDNA samplers will accelerate the application of eDNA-based methods in parallel with other surveying techniques (environmental and geophysical). Automated eDNA samplers are already being used in marine environments, attached to a fixed point for monitoring over time (Mynott 2019), with the ROVs (Everett and Park 2018) or coupling an environmental sample processor with an AUV (Yamahara et al. 2019a). Automated environmental sampling has also been applied to freshwater monitoring (Searcy et al. 2022).

For industrial applications, the development of new autonomous sampling equipment will reduce offshore vessel and staff time, particularly if deployed from infrastructure rather than from a vessel (which is frequently required for ROV/AUV deployment). The lower survey costs from decreased vessel time may allow for enhanced environmental monitoring if the saving is used to collect more samples (McLean et al. 2020). The same ROV/AUV used to collect eDNA can also be used to obtain conventional biological samples as well as other types of data.

Automated and drone-based sampling of freshwater and soil samples can also be considered as part of environmental surveying (Robinson et al. 2022). The application of drone technology has been relatively limited to date (Doi et al. 2017), although the weight of samples or the drone technology do not limit its use. The advantage of this approach would be that remote sensing could obtain broad environmental data for an area, followed by subsequent individually selected eDNA sampling stations for soil and water, to calibrate the habitat data. This would significantly reduce the time and effort of field ecologists.

7.2.2 In-Field Analysis

Mobile laboratory technologies already exist, allowing for eDNA samples to be processed from extraction to sequencing in the field within marine and freshwater environments (Chang et al. 2020; Hansen et al. 2020; Urban et al. 2021). However, further developments are needed to improve efficiency and adjust these mobile technologies to regulatory standards for widespread application (Environment Agency 2021). If made widely available to industry, automated sampling and sample processing of eDNA could be deployed on buoys and be used together with other systems used for passive acoustic monitoring for marine mammals (Kowarski et al. 2020) or plankton sampling in the marine environment (e.g. Pitois et al. 2021).

In-field methods for rapid single-species detection are also under development, including methods using CRISPR (M. Williams et al. 2019) and various other isothermal amplification methods (Fast et al. 2020). Systematic application of these tools in the field requires investment in the development of reliable methods for simple, fast, in-field extraction of high-quality DNA in a contamination-secure manner. This will reduce the time between sampling and obtaining results from days to hours or even minutes. This makes such methods ideal for rapid detection of INNS, harmful taxa or pollution indicator species. However, this is unlikely to replace visual and PAM observations for marine mammals and reptiles, where a

response is required immediately after the observation. For example, a visual observation might lead to immediately stopping high-decibel activity at offshore sites.

7.2.3 Other Substrates for DNA Sampling

Sampling eDNA from air is in its infancy compared to sampling eDNA from water or soil. However, over the last few years, there have been an increasing number of studies using airborne DNA for assessing plant communities (M. D. Johnson et al. 2021; Banchi et al. 2020), fungal communities (Rosa et al. 2020), vertebrates (Lynggaard et al. 2022), invertebrates (Roger et al. 2022), and eukaryotes (Aalismail et al. 2021). Analysing DNA from the air offers a totally new approach to assess biodiversity in terrestrial environments. Its future use for baselining surveys for industrial projects and their environmental impact assessment should be considered, but some further methodological development may be necessary before the approach is mature and ready to use.

7.3 Laboratory Work and Analysis

These components are primarily addressed in JIP-34 Chapters 3 and 4. They are addressed briefly here from a perspective of how the future eDNA-based methods can deal with the current challenges of data generation and analysis.

7.3.1 Technological Innovations

Tools and technologies have rapidly developed since the first proposed use of environmental DNA in biodiversity monitoring, and they will continue to do so. The high-throughput sequencing platforms used in many environmental genomic approaches are continuously developing - with new instruments, producing more sequences more rapidly (e.g. Illumina NovaSeq) or potentially enabling the portable real-time DNA/RNA sequencing at an affordable price (Oxford Nanopore MinION). The development of nanopore-like technology is of particular interest for biomonitoring as it responds to a need for a rapid and cost-effective tool to generate sequencing data. Moreover, the long reads produced by nanopore technology enable more accurate recovery of taxonomic and phylogenetic diversity than short amplicon sequencing (Kreherwinkel et al. 2019). Although the adoption of this nanopore technology has long been held back by technical issues, many previous limitations to the technical feasibility of these instruments continue to be improved on, and their potential application to biodiversity monitoring and ecosystem health assessment is being recognized (Urban et al. 2021).

7.3.2 Taxonomic Reference Database

Building a comprehensive reference database is another important challenge for the future of eDNA-based monitoring. The accuracy of biodiversity surveys depends on whether the species can be correctly identified. This is particularly important when environmental genomics methods are proposed to replace the morphology-based identifications. Unfortunately, many morphospecies, especially among the invertebrates and microbial eukaryotes used as bioindicators, do not have reference sequences in public databases. There are ongoing efforts to fill the gaps in the current DNA barcode databases (e.g. Diat.barcode, Rimet et al. 2019), but, these efforts usually focus on one marker that has been selected based on current research, which limits the possibility to change the marker if a more effective marker is found. It is expected that the future development of reference databases will use long-read sequencing technologies (e.g. PacBio) to sequence whole genomes (e.g. mitochondrial genomes) or entire operons (e.g. rRNA genes) which will provide better taxonomic resolution and enable selection of more convenient markers for a given taxon (Jamy et al. 2020).

7.3.3 Species Detection

The reference database is essential for designing better primers for metabarcoding and single-species detection. New primers can largely increase the proportion of target taxa within the metabarcoding datasets, as demonstrated with marine vertebrates (Valsecchi et al. 2020) and freshwater aquatic invertebrates (Brantschen et al. 2022). Optimizing primers and probes is particularly important for the development of single species assays. This will likely increase the use of eDNA-based methods if their detection capability is improved for the target species, especially with regards to INNS, protected and indicator taxa.

7.3.4 Abundance Measures

Future eDNA-based biomonitoring will also need to overcome the limitations related to the semi-quantitative nature of DNA sequence data (Luo et al. 2022). In most cases, there is no direct relationship between the number of specimens and number of sequences, which impedes the comparison of eDNA and conventional abundance data. Calibration against conventional data can be used to correct biases between eDNA concentration and biomass or abundance, although this should consider a range of environmental variables that affect eDNA decay rate as well as the variable biological relationship between shedding rate and biomass or abundance. However, all survey methods, including conventional ones, have biases that should also be considered. Recently, several studies proposed to correct different biases that impact the relationship between biomass, eDNA concentration and the number of sequences through:

- the use of multiple markers with different target regions and biases (Bucklin et al. 2021).
- including an internal control of known concentration of DNA, or “spike-in”, in samples (Stoeckle, Ausubel, and Coogan 2022).
- sequencing mock samples that include known concentrations of target species (McLaren, Willis, and Callahan 2019; Shelton et al. 2022)
- sequencing the same sample that has been amplified using different number of cycles during PCR (Silverman et al. 2021).

Routine use of these methods, while more expensive, will result in better quantification of each species' eDNA concentration (Gold et al. 2023). This may ultimately result in eDNA-based methods being used to better estimate biomass data, reducing the need to collect as many conventional samples to obtain population-level data (Jungbluth et al. 2022; Di Muri et al. 2020).

7.3.5 Consistency of Results

The automation of eDNA analysis is burgeoning, including sample processing, bioinformatics and data analysis. The use of robotics will minimise risk of any subjectivity or human errors and enhance consistency of results (Tegally et al. 2020). At the same time, the continued development of bioinformatic tools is expected to resolve problems related to the interpretation of sequence data, particularly to improve the taxonomic assignment of sequences and to correct technical errors. The complementarity of eDNA-based and conventional methods requires better fit between molecular and morphological identifications, which remains a challenge for the future developments of eDNA-based biomonitoring.

7.4 Metrics of Ecosystem Health

To date, most conventional indices of ecological quality status are based on the presence and abundance of species and using these to generate a score. Only species that are

assigned to an ecological category or an indicator value are included in the calculation of biotic indices (Pawlowski et al. 2018).

Future eDNA-based metrics will overcome this limitation by promoting a more holistic approach to ecosystem condition assessment using a taxonomy-free approach. This can be done either by assigning indicator values directly to sequence data (Apothéoz-Perret-Gentil et al. 2017) or by using machine learning to create models based on the full metabarcoding dataset, without any taxonomic criteria (Cordier et al. 2020). These taxonomy-free models work by learning the biological community signature that defines a healthy ecosystem and predicting how that signature changes as the ecosystem degrades or improves. Once trained, the model can then assign unknown samples to an ecosystem health class or a position on the degradation gradient. Machine learning models have been used to predict the impacts of marine aquaculture (Cordier et al. 2017; 2018; Frühe et al. 2020), as well as to assess the ecological status of coastal marine habitats (DiBattista et al. 2020), and rivers using diatoms (Feio et al. 2020) and phytoplankton (Fan et al. 2020).

The machine learning approach has been shown to be more powerful for ecosystem assessment than taxonomy-based indices (Cordier et al. 2018), but it is not yet clear how transferable these models are between habitats and locations. Future developments in this direction will require much more extensive training datasets that will take a broad range of environmental parameters, including seasonality, biogeography, etc., into consideration. Building a comprehensive database of DNA metabarcodes associated with industrial impacts is essential to develop novel metrics for eDNA-based monitoring.

7.5 Standardisation

To improve management and conservation efforts and enhance the implementation of eDNA-based monitoring, more thorough guidelines, or standardisation of environmental genomics workflows is needed. The eDNA sampling process is discussed in greater detail in IOGP JIP34, Chapter 2 and the laboratory and bioinformatics components are discussed by the IOGP JIP34 Chapters 3 and 4 respectively (e.g. MIQE guidelines for reporting information on qPCR experiments (Bustin et al. 2009)). Here, we will report the standardisation efforts that have already been done in this field and highlight their further development.

Governmental and intergovernmental regulators consider the relative merits and calibration stages of eDNA-based methods against conventional methods prior to recommending them as a monitoring method (Aylagas et al. 2020; Bruce et al. 2021; Hering et al. 2018). Therefore, it is of critical importance to promote a formal acceptance of eDNA-based methods. This can be done by publishing eDNA guidelines supported by governmental agencies, as it has been done in Japan, Canada and Switzerland (The eDNA Society 2019; Pawlowski et al. 2020; Abbott et al. 2021). Recent initiatives to incorporate eDNA methods into regulatory monitoring and acceptance in legal national and international instruments include:

- a technical report for the routine sampling of benthic diatoms from rivers and lakes adapted for metabarcoding analyses, published by the European Committee for Standardization (FD CEN/TR17245:2018)
- publication of a protocol on metagenomic analysis of meiofauna in marine environments by the International Standardization Organization (ISO 23732:2021(E))
- creation of a working group on DNA and eDNA methods (CEN/TC230/WG28) and publication of the first standard on “Collecting and preserving samples for capture of environmental DNA in aquatic environments from water samples” in 2023 (SN EN 17805:23).

Still, these efforts are at a very early stage. Most of the research on eDNA-based biomonitoring present in this report and elsewhere remains at the academic level. To be routinely applicable, the environmental genomic methods used in this field need to go through a long process of standardisation and optimization. This process has already begun and will certainly continue in the future.

7.6 Summary

To conclude, the future of eDNA-based biomonitoring depends on three main factors: technological advances, scientific evidence and regulatory acceptance. Technology is advancing rapidly, offering a vision of future biomonitoring which relies on automated eDNA samplers and robotics which generate a continuous flow of sequence data that are analysed using machine learning or other AI tools. However, this vision might not be easily accepted by those practicing conventional biomonitoring. For many, eDNA-based methods should help to detect and identify species but not necessarily to change the way the ecological status is assessed and the bioindicator used for that assessment. eDNA-based methods should continue to be calibrated against conventional methods to ensure that results are comparable under a range of conditions allowing interoperability between different datasets. Scientific evidence is needed to demonstrate the accuracy of eDNA-based technology, but its implementation cannot take place without regulatory acceptance. Hence, the combination of these three factors is essential for the success of eDNA-based monitoring tools.

As shown throughout this report, there is a consensus that eDNA-based methods should be integrated into future biomonitoring studies. Whether this will be done by replacing or in combination with conventional methods will depend on the habitat, target taxa, data required and survey objectives. It should be noted that many new eDNA-based methods are still at the level of academic research. Further development and optimization may be required before these methods can be used in routine biomonitoring. However, the benefits of using eDNA will certainly outweigh these efforts, making future biomonitoring of industrial operations more efficient and beneficial for ecosystem health and biodiversity conservation.

Glossary

Assay	Analysis used for DNA samples, often referring to PCR reaction conditions and primers, but may also be used to encompass different molecular biology protocols in a workflow
ASV	Amplicon Sequence Variant – metabarcoding datasets of individual high-quality sequences because sequences that contain errors are typically filtered out using denoising algorithms. Often used as an alternative or in combination with clustering (OTUs) for taxon delimitation.
AUV	Autonomous Underwater Vehicle
Benthic	Organisms living at the bottom of a water body (seafloor, river or lake bottom)
Bioinformatics	The application of genomics, computational biology, statistics, and programming in which DNA sequences or other biological data are processed, analysed and integrated for research or communications.
Bulk sample	Sample composed of unsorted organisms, mainly referring to invertebrates
Conventional	For the purposes of this document, any survey method that is not DNA-based
COI	Mitochondrial gene (Cytochrome C Oxidase <i>I</i>) commonly used for animal DNA barcoding. For metabarcoding, it is frequently used for analysis of invertebrate taxa.
DEM	Digital Elevation Model – model of topography
Diatom	Single-celled algae (typically with a silica frustule) frequently used for water quality assessments.
DSM	Digital Surface Model – DEM with any additional structures (vegetation, anthropogenic)
eDNA	Short for ‘environmental DNA’. Refers to DNA in the environment, including DNA released from organisms through excretion, shedding, mucous secretions, saliva etc. This can be collected in environmental samples (e.g. water, sediment) and used to identify the organisms that it originated from. eDNA in water is broken down by environmental processes over a period of days to weeks. It can travel some distance from the point at which it was released from the organism, particularly in running water. eDNA is sampled in low concentrations and can be degraded (i.e. broken into short fragments), which limits the analysis options.
Elasmobranch	Taxonomic group of cartilaginous fish (sharks, rays).
Eukaryote	Organism with a nucleus (animals, plants, fungi, algae).
GCN	Great Crested Newt
iDNA	Invertebrate-derived DNA; DNA sourced from invertebrates (e.g. mosquitoes and leeches) which is from target organisms (e.g. vertebrates) that the insects feed on.
INNS	Invasive and Non-Native Species
Malaise trap	Trap commonly used to sample flying insects
Metabarcoding	Refers to the simultaneous identification of multiple species from DNA using barcode genes. Target sections of DNA are amplified through PCR, followed by high-throughput sequencing and bioinformatics processing. Can identify hundreds of species in each sample, and 100+ different samples can be processed in parallel to reduce cost.
Metagenomics	Non-targeted sequencing of DNA within a mixed sample compared to metabarcoding which sequences a specific gene region (e.g. Bista et al. 2018)

Metatranscriptomics	Study of the transcriptome, the sequences of RNA within a mixed sample. Also used to identify the “active” community (e.g. Knapik et al. 2020).
Morphology/Morphological analysis	Assessment of organisms within a sample based on visible physical characteristics (commonly used identification).
-Omics	Colloquialism used to encompass transcriptomics, genomics, metabolomics etc.
OTU	Operational Taxonomic Unit – sequences clustered above a given similarity threshold to assign quasi-species status.
PAM	Passive Acoustic Monitoring – bio-acoustic surveying
Passive sampler	eDNA sampler without any moving (active) components, typically using a material that adsorbs eDNA.
PCR	Polymerase Chain Reaction: a process by which millions of copies of a particular DNA segment are produced through a series of heating and cooling steps, known as an ‘amplification’ process. One of the most common processes in molecular biology and a precursor to most sequencing-based analyses.
dPCR	Digital PCR - This method is similar to qPCR in application but is more robust to inhibitors, does not require standards, and offers greater precision for quantifying DNA concentration (e.g. Baker et al. 2018). Encompasses droplet digital PCR (ddPCR)
qPCR	Quantitative PCR - A fluorescent signal is emitted as DNA is amplified which allows DNA concentration to be quantified; thus, this approach can be used to quantify the concentration of targeted DNA.
Pelagic	Adjective for describing biology in the marine water column
PLFA	Phospholipid fatty acid – conventional analysis used for assessing microbial communities
Pitfall trap	Trap used to sample surface-dwelling invertebrates
Primer	Short sections of synthesised DNA that bind to either end of the DNA segment to be amplified by PCR. Can be designed to be totally specific to a particular species (so that only that species’ DNA will be amplified from a community DNA sample), or to be very general so that a wide range of species’ DNA will be amplified. Good design of primers is one of the critical factors in DNA-based monitoring.
Prokaryote	Organism without nucleus (bacteria, archaea)
PSD	Particle Size Distribution – the weight distribution of different grain sizes (mud, sand, gravel) of a sediment
Reference database	Publicly available database of DNA sequences with taxonomic assignments from many species around the world. These databases serve as a reference against which unknown sequences can be queried and identified to a taxon. The most accessed database is NCBI, which is maintained by the US National Institute of Health. Anyone can search for DNA sequences at https://www.ncbi.nlm.nih.gov .
ROV	Remotely Operated Vehicle
(DNA) Sequence	A section of DNA. A DNA sequence is made up of four nucleotide bases represented by the letters A, T, C & G. The precise order of these letters is used to compare genetic similarity among individuals or species and to identify species using reference databases. In high-throughput sequencing analyses (e.g. DNA metabarcoding), identical copies of the same sequence are obtained for each species in the sample. The number of copies

Spike-in	obtained per species is known as the number of sequencing reads, and this can be related to the relative abundance of species. Addition of DNA to a sample. This artificially added DNA can be used to assess and correct for biases in PCR and bioinformatics pipelines.
Thermocline	A water layer within which there is a rapid change in temperature (typically marine and lakes).
18S (rRNA)	Gene coding for small subunit ribosomal RNA. Commonly used in metabarcoding for targeting eukaryotes/metazoans.



Appendix%201%20R
eview%20Database.xls

References

- Aalismail, Nojood A., Rubén Díaz-Rúa, Nathan Geraldi, Michael Cusack, and Carlos M. Duarte. 2021. "Diversity and Sources of Airborne Eukaryotic Communities (AEC) in the Global Dust Belt over the Red Sea." *Earth Systems and Environment* 5 (2): 459–71. <https://doi.org/10.1007/s41748-021-00219-4>.
- Abbott, Cathryn, Mark Coulson, Nellie Gagné, Anaïs Lacoursière, Robert Bajno, Charise Dietrich, and Shannan May-McNally. 2021. "Guidance on the Use of Targeted Environmental DNA (eDNA) Analysis for the Management of Aquatic Invasive Species and Species at Risk." *DFO Canadian Science Advisory Secretariat Research Document*.
- Abrams, Jesse F., Lisa A. Hörig, Robert Brozovic, Jan Axtner, Alex Crampton-Platt, Azlan Mohamed, Seth T. Wong, Rahel Sollmann, Douglas W. Yu, and Andreas Wilting. 2019. "Shifting up a Gear with iDNA : From Mammal Detection Events to Standardised Surveys." Edited by Helen Wheeler. *Journal of Applied Ecology* 56 (7): 1637–48. <https://doi.org/10.1111/1365-2664.13411>.
- Adams, Clare, Luke Hoekstra, Morgan Muell, and Fredric Janzen. 2019. "A Brief Review of Non-Avian Reptile Environmental DNA (eDNA), with a Case Study of Painted Turtle (*Chrysemys Picta*) eDNA Under Field Conditions." *Diversity* 11 (4): 50. <https://doi.org/10.3390/d11040050>.
- Alexander, Jason B., Michael J. Marnane, Travis S. Elsdon, Michael Bunce, Paweena Sitaworawet, Se Songploy, Sarin Chaiyakul, and Euan S. Harvey. 2023. "Using Environmental DNA to Better Inform Decision Making around Decommissioning Alternatives for Offshore Oil and Gas Infrastructure." *Science of The Total Environment* 901 (November): 165991. <https://doi.org/10.1016/j.scitotenv.2023.165991>.
- Alexander, Jason B, Michael J Marnane, Travis S Elsdon, Michael Bunce, Se Songploy, Paweena Sitaworawet, and Euan S. Harvey. 2022. "Complementary Molecular and Visual Sampling of Fish on Oil and Gas Platforms Provides Superior Biodiversity Characterisation." *Marine Environmental Research* 179: 11.
- Alexander, Jason B., Michael J. Marnane, Justin I. McDonald, Sherralee S. Lukehurst, Travis S. Elsdon, Tiffany Simpson, Shawn Hinz, Michael Bunce, and Euan S. Harvey. 2023. "Comparing Environmental DNA Collection Methods for Sampling Community Composition on Marine Infrastructure." *Estuarine, Coastal and Shelf Science* 283 (April): 108283. <https://doi.org/10.1016/j.ecss.2023.108283>.
- Allen, Michael C., Robert Kwait, Anthony Vastano, Alex Kisurin, Isabelle Zoccolo, Benjamin D. Jaffe, Jordan C. Angle, Brooke Maslo, and Julie L. Lockwood. 2023. "Sampling Environmental DNA from Trees and Soil to Detect Cryptic Arboreal Mammals." *Scientific Reports* 13 (1): 180. <https://doi.org/10.1038/s41598-023-27512-8>.
- Alve, Elisabeth, Sergei Korsun, Joachim Schönfeld, Noortje Dijkstra, Elena Golikova, Silvia Hess, Katrine Husum, and Giuliana Panieri. 2016. "Foram-AMBI: A Sensitivity Index Based on Benthic Foraminiferal Faunas from North-East Atlantic and Arctic Fjords, Continental Shelves and Slopes." *Marine Micropaleontology* 122 (January): 1–12. <https://doi.org/10.1016/j.marmicro.2015.11.001>.
- Anderson, James T., Florencia L. Zilli, Luciana Montalto, Mercedes R. Marchese, Matthew McKinney, and Yong-Lak Park. 2013. "Sampling and Processing Aquatic and Terrestrial Invertebrates in Wetlands." In *Wetland Techniques*, edited by James T. Anderson and Craig A. Davis, 143–95. Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-007-6931-1_5.
- Ankley, Phillip J., Yuwei Xie, Tyler A. Black, Abigail DeBofsky, McKenzie Perry, Michael J. Paterson, Mark Hanson, Scott Higgins, John P. Giesy, and Vince Palace. 2021. "Using Zooplankton Metabarcoding to Assess the Efficacy of Different Techniques to

- Clean-up an Oil-Spill in a Boreal Lake." *Aquatic Toxicology* 236 (July): 105847. <https://doi.org/10.1016/j.aquatox.2021.105847>.
- Ankley, Phillip J., Yuwei Xie, Sonya Havens, Lisa Peters, Lauren Timlick, Jose Luis Rodriguez-Gil, John P. Giesy, and Vince P. Palace. 2022. "RNA Metabarcoding Helps Reveal Zooplankton Community Response to Environmental Stressors." *Environmental Pollution* 292 (January): 118446. <https://doi.org/10.1016/j.envpol.2021.118446>.
- Apothéloz-Perret-Gentil, Laure, Agnès Bouchez, Tristan Cordier, Arielle Cordonier, Julie Guéguen, Frederic Rimet, Valentin Vasselon, and Jan Pawlowski. 2021. "Monitoring the Ecological Status of Rivers with Diatom eDNA Metabarcoding: A Comparison of Taxonomic Markers and Analytical Approaches for the Inference of a Molecular Diatom Index." *Molecular Ecology* 30 (13): 2959–68. <https://doi.org/10.1111/mec.15646>.
- Apothéloz-Perret-Gentil, Laure, Arielle Cordonier, François Straub, Jennifer Iseli, Philippe Esling, and Jan Pawlowski. 2017. "Taxonomy-Free Molecular Diatom Index for High-Throughput eDNA Biomonitoring." *Molecular Ecology Resources* 17 (6): 1231–42. <https://doi.org/10.1111/1755-0998.12668>.
- Appel, Giulliana, Ubirajara D. Capaverde, Leonardo Queiroz De Oliveira, Lucas G. Do Amaral Pereira, Valéria Da Cunha Tavares, Adrià López-Baucells, William E. Magnusson, Fabrício Beggiato Baccaro, and Paulo E. D. Bobrowiec. 2022. "Use of Complementary Methods to Sample Bats in the Amazon." *Acta Chiropterologica* 23 (2). <https://doi.org/10.3161/15081109ACC2021.23.2.017>.
- Armbruster, Melanie, Tim Goodall, Penny R. Hirsch, Nick Ostle, Jeremy Puissant, Kate C. Fagan, Richard F. Pywell, and Robert I. Griffiths. 2021. "Bacterial and Archaeal Taxa Are Reliable Indicators of Soil Restoration across Distributed Calcareous Grasslands." *European Journal of Soil Science* 72 (6): 2430–44. <https://doi.org/10.1111/ejss.12977>.
- Aylagas, Eva, Ángel Borja, Iñigo Muxika, and Naiara Rodríguez-Ezpeleta. 2018. "Adapting Metabarcoding-Based Benthic Biomonitoring into Routine Marine Ecological Status Assessment Networks." *Ecological Indicators* 95 (December): 194–202. <https://doi.org/10.1016/j.ecolind.2018.07.044>.
- Aylagas, Eva, Angel Borja, Xavier Pochon, Anastasija Zaiko, Nigel Keeley, Kat Bruce, Peiying Hong, et al. 2020. "Translational Molecular Ecology in Practice: Linking DNA-Based Methods to Actionable Marine Environmental Management." *Science of The Total Environment* 744 (November): 140780. <https://doi.org/10.1016/j.scitotenv.2020.140780>.
- Aylagas, Eva, Iñaki Mendibil, Ángel Borja, and Naiara Rodríguez-Ezpeleta. 2016. "Marine Sediment Sample Pre-Processing for Macroinvertebrates Metabarcoding: Mechanical Enrichment and Homogenization." *Frontiers in Marine Science* 3 (October). <https://doi.org/10.3389/fmars.2016.00203>.
- Bahnmann, Barbara, Tereza Mašínová, Rune Halvorsen, Marie L. Davey, Petr Sedlák, Michal Tomšovský, and Petr Baldrian. 2018. "Effects of Oak, Beech and Spruce on the Distribution and Community Structure of Fungi in Litter and Soils across a Temperate Forest." *Soil Biology and Biochemistry* 119 (April): 162–73. <https://doi.org/10.1016/j.soilbio.2018.01.021>.
- Baillet, Bonnie, Agnes Bouchez, Alain Franc, Jean-Marc Frigerio, François Keck, Satu-Maaria Karjalainen, Frederic Rimet, Susanne Schneider, and Maria Kahlert. 2019. "Molecular versus Morphological Data for Benthic Diatoms Biomonitoring in Northern Europe Freshwater and Consequences for Ecological Status." *Metabarcoding and Metagenomics* 3 (June): e34002. <https://doi.org/10.3897/mbmg.3.34002>.

- Baker, C. Scott, Debbie Steel, Sharon Nieu Kirk, and Holger Klinck. 2018. "Environmental DNA (eDNA) From the Wake of the Whales: Droplet Digital PCR for Detection and Species Identification." *Frontiers in Marine Science* 5 (April): 133. <https://doi.org/10.3389/fmars.2018.00133>.
- Bálint, Miklós, Carsten Nowak, Orsolya Marton, Steffen U. Pauls, Claudia Wittwer, José Luis Aramayo, Arne Schulze, Thierry Chambert, Bernardino Cocchiararo, and Martin Jansen. 2018. "Accuracy, Limitations and Cost-Efficiency of eDNA-Based Community Survey in Tropical Frogs." *Molecular Ecology Resources* 18 (6): 43.
- Banchi, Elisa, Claudio G. Ametrano, Enrico Tordoni, David Stanković, Silvia Ongaro, Mauro Tretlach, Alberto Pallavicini, et al. 2020. "Environmental DNA Assessment of Airborne Plant and Fungal Seasonal Diversity." *Science of The Total Environment* 738 (October): 140249. <https://doi.org/10.1016/j.scitotenv.2020.140249>.
- Barrenechea Angeles, Ines, Maria Lorena Romero-Martínez, Marco Cavaliere, Stefano Varrella, Fabio Francescangeli, Roberta Piredda, Maria Grazia Mazzocchi, et al. 2023. "Encapsulated in Sediments: eDNA Deciphers the Ecosystem History of One of the Most Polluted European Marine Sites." *Environment International* 172 (February): 107738. <https://doi.org/10.1016/j.envint.2023.107738>.
- Basset, Yves, Mehrdad Hajibabaei, Michael T. G. Wright, Anakena M. Castillo, David A. Donoso, Simon T. Segar, Daniel Souto-Vilarós, et al. 2022. "Comparison of Traditional and DNA Metabarcoding Samples for Monitoring Tropical Soil Arthropods (Formicidae, Collembola and Isoptera)." *Scientific Reports* 12 (1): 10762. <https://doi.org/10.1038/s41598-022-14915-2>.
- Bessey, Cindy, Yuan Gao, Yen Bach Truong, Haylea Miller, Simon Neil Jarman, and Oliver Berry. 2021. "Comparison of Materials for Rapid Passive Collection of Environmental DNA." *bioRxiv Preprint*, November. <https://doi.org/10.1101/2021.11.08.467842>.
- Besson, Marc, Jamie Alison, Kim Bjerge, Thomas Gorochofski, Toke Høye, Tommaso Jucker, Hjalte Mann, and Christopher Clements. 2022. "Towards the Fully Automated Monitoring of Ecological Communities." Preprint. Preprints. <https://doi.org/10.22541/au.165063792.29640103/v1>.
- Bienert, Friederike, Sébastien De Danieli, Christian Miquel, Eric Coissac, Carole Poillot, Jean-Jacques Brun, and Pierre Taberlet. 2012. "Tracking Earthworm Communities from Soil DNA." *Molecular Ecology* 21 (8): 2017–30. <https://doi.org/10.1111/j.1365-294X.2011.05407.x>.
- Biggs, Jeremy, Naomi Ewald, Alice Valentini, Coline Gaboriaud, Richard A. Griffiths, Jim Foster, John W. Wilkinson, A Arnett, Penny Williams, and Francesca Dunn. 2014. "Analytical and Methodological Development for Improved Surveillance of the Great Crested Newt - WC1067." Defra Project WC1067. Freshwater Habitats Trust:Oxford.
- Biggs, Jeremy, Naomi Ewald, Alice Valentini, Coline Gaboriaud, Tony Dejean, Richard A. Griffiths, Jim Foster, et al. 2015. "Using eDNA to Develop a National Citizen Science-Based Monitoring Programme for the Great Crested Newt (*Triturus cristatus*)." *Biological Conservation* 183: 19–28. <https://doi.org/10.1016/j.biocon.2014.11.029>.
- Bik, Holly M., Kenneth M. Halanych, Jyotsna Sharma, and W. Kelley Thomas. 2012. "Dramatic Shifts in Benthic Microbial Eukaryote Communities Following the Deepwater Horizon Oil Spill." Edited by Ahmed Moustafa. *PLoS ONE* 7 (6): e38550. <https://doi.org/10.1371/journal.pone.0038550>.
- Blabolil, Petr, Nathan P. Griffiths, Bernd Hänfling, Tomáš Jůza, Vladislav Draštík, Jelena Knežević-Jarić, Romulo dos Santos, Tomáš Mrkvička, and Jiří Peterka. 2022. "The True Picture of Environmental DNA, a Case Study in Harvested Fishponds." *Ecological Indicators* 142 (September): 109241. <https://doi.org/10.1016/j.ecolind.2022.109241>.

- Blackman, Rosetta C., Jeanine Brantschen, Jean-Claude Walser, Remo Wüthrich, and Florian Altermatt. 2021. "Monitoring Invasive Alien Macroinvertebrate Species with Environmental DNA." *River Research and Applications* 38 (8): 1400–1412. <https://doi.org/10.1002/rra.3947>.
- Blackman, Rosetta C., Kar Keun Sean Ling, Lynsey R. Harper, Peter Shum, Bernd Hänfling, and Lori Lawson-Handley. 2020. "Targeted and Passive Environmental DNA Approaches Outperform Established Methods for Detection of Quagga Mussels, *Dreissena Rostriformis Bugensis* in Flowing Water." *Ecology and Evolution* 10 (23): 13248–59. <https://doi.org/10.1002/ece3.6921>.
- Böckmann, Elias, Alexander Pfaff, Michael Schirrmann, and Michael Pflanz. 2021. "Rapid and Low-Cost Insect Detection for Analysing Species Trapped on Yellow Sticky Traps." *Scientific Reports* 11 (1): 10419. <https://doi.org/10.1038/s41598-021-89930-w>.
- Boetzi, Fabian A., Elena Ries, Gudrun Schneider, and Jochen Krauss. 2018. "It's a Matter of Design—How Pitfall Trap Design Affects Trap Samples and Possible Predictions." *PeerJ* 6 (June): e5078. <https://doi.org/10.7717/peerj.5078>.
- Boivin-Delisle, Damien, Martin Laporte, Frédéric Burton, René Dion, Eric Normandeau, and Louis Bernatchez. 2021. "Using Environmental DNA for Biomonitoring of Freshwater Fish Communities: Comparison with Established Gillnet Surveys in a Boreal Hydroelectric Impoundment." *Environmental DNA* 3 (1): 105–20. <https://doi.org/10.1002/edn3.135>.
- Bongers, Tom. 1990. "The Maturity Index: An Ecological Measure of Environmental Disturbance Based on Nematode Species Composition." *Oecologia* 83 (1): 14–19. <https://doi.org/10.1007/BF00324627>.
- Booth, Cormac G., Rachael R. Sinclair, and John Harwood. 2020. "Methods for Monitoring for the Population Consequences of Disturbance in Marine Mammals: A Review." *Frontiers in Marine Science* 7 (February): 115. <https://doi.org/10.3389/fmars.2020.00115>.
- Borja, A, J Franco, and V Pérez. 2000. "A Marine Biotic Index to Establish the Ecological Quality of Soft-Bottom Benthos Within European Estuarine and Coastal Environments." *Marine Pollution Bulletin* 40 (12): 1100–1114. [https://doi.org/10.1016/S0025-326X\(00\)00061-8](https://doi.org/10.1016/S0025-326X(00)00061-8).
- Borrego-Ramos, María, Eloy Bécares, Pedro García, Alejandro Nistal, and Saúl Blanco. 2021. "Epiphytic Diatom-Based Biomonitoring in Mediterranean Ponds: Traditional Microscopy versus Metabarcoding Approaches." *Water* 13 (10): 1351. <https://doi.org/10.3390/w13101351>.
- Boussarie, Germain, Judith Bakker, Owen S. Wangensteen, Stefano Mariani, Lucas Bonnin, Jean-Baptiste Juhel, Jeremy J. Kiszka, et al. 2018. "Environmental DNA Illuminates the Dark Diversity of Sharks." *Science Advances* 4 (5): eaap9661. <https://doi.org/10.1126/sciadv.aap9661>.
- Bradley, Douglas L., Kevin C. Morey, Danielle A. Bourque, Brooks Fost, Tzitziki Loeza-Quintana, and Robert H. Hanner. 2022. "Environmental DNA Detection and Abundance Estimates Comparable to Conventional Methods for Three Freshwater Larval Species at a Power Plant Discharge." *Environmental DNA*, February, edn3.286. <https://doi.org/10.1002/edn3.286>.
- Brakstad, Odd G., Mimmi Throne-Holst, Roman Netzer, Donald M. Stoeckel, and Ronald M. Atlas. 2015. "Microbial Communities Related to Biodegradation of Dispersed M Acondo Oil at Low Seawater Temperature with Norwegian Coastal Seawater." *Microbial Biotechnology* 8 (6): 989–98. <https://doi.org/10.1111/1751-7915.12303>.
- Brannock, Pamela M., Jyotsna Sharma, Holly M. Bik, W. Kelley Thomas, and Kenneth M. Halanych. 2017. "Spatial and Temporal Variation of Intertidal Nematodes in the

- Northern Gulf of Mexico after the Deepwater Horizon Oil Spill." *Marine Environmental Research* 130 (September): 200–212.
<https://doi.org/10.1016/j.marenvres.2017.07.008>.
- Brannock, Pamela M., Damien S. Waits, Jyotsna Sharma, and Kenneth M. Halanych. 2014. "High-Throughput Sequencing Characterizes Intertidal Meiofaunal Communities in Northern Gulf of Mexico (Dauphin Island and Mobile Bay, Alabama)." *The Biological Bulletin* 227 (2): 161–74. <https://doi.org/10.1086/BBLv227n2p161>.
- Brantschen, Jeanine, Rosetta C. Blackman, Jean-Claude Walser, and Florian Altermatt. 2021. "Environmental DNA Gives Comparable Results to Morphology-Based Indices of Macroinvertebrates in a Large-Scale Ecological Assessment." Edited by Hideyuki Doi. *PLOS ONE* 16 (9): e0257510. <https://doi.org/10.1371/journal.pone.0257510>.
- Brantschen, Jeanine, Loïc Pellissier, Jean-Claude Walser, and Florian Altermatt. 2022. "Evaluation of Primer Pairs for eDNA-based Assessment of Ephemeroptera, Plecoptera, and Trichoptera across a Biogeographically Diverse Region." *Environmental DNA*, August, edn3.342. <https://doi.org/10.1002/edn3.342>.
- Bruce, Kat, Rosetta Blackman, Sarah J. Bourlat, Ann Micaela Hellström, Judith Bakker, Iliana Bista, Kristine Bohmann, et al. 2021. *A Practical Guide to DNA-Based Methods for Biodiversity Assessment*. Pensoft Publishers. <https://doi.org/10.3897/ab.e68634>.
- Brys, Rein, Annelies Haegeman, David Halfmaerten, Sabrina Neyrinck, Ariane Staelens, Johan Auwerx, and Tom Ruttink. 2021. "Monitoring of Spatiotemporal Occupancy Patterns of Fish and Amphibian Species in a Lentic Aquatic System Using Environmental DNA." *Molecular Ecology* 30 (13): 3097–3110.
<https://doi.org/10.1111/mec.15742>.
- Bucklin, Ann, Katja T. C. A. Peijnenburg, Ksenia N. Kosobokova, Todd D. O'Brien, Leocadio Blanco-Bercial, Astrid Cornils, Tone Falkenhaug, et al. 2021. "Toward a Global Reference Database of COI Barcodes for Marine Zooplankton." *Marine Biology* 168 (6): 78. <https://doi.org/10.1007/s00227-021-03887-y>.
- Burton, A. Cole, Eric Neilson, Dario Moreira, Andrew Ladle, Robin Steenweg, Jason T. Fisher, Erin Bayne, and Stan Boutin. 2015. "Wildlife Camera Trapping: A Review and Recommendations for Linking Surveys to Ecological Processes." Edited by Phil Stephens. *Journal of Applied Ecology* 52 (3): 675–85. <https://doi.org/10.1111/1365-2664.12432>.
- Bustin, Stephen A, Vladimir Benes, Jeremy A Garson, Jan Hellemans, Jim Huggett, Mikael Kubista, Reinhold Mueller, et al. 2009. "The MIQE Guidelines: Minimum Information for Publication of Quantitative Real-Time PCR Experiments." *Clinical Chemistry* 55 (4): 611–22. <https://doi.org/10.1373/clinchem.2008.112797>.
- Cahill, Abigail E., John K. Pearman, Angel Borja, Laura Carugati, Susana Carvalho, Roberto Danovaro, Sarah Dashfield, et al. 2018. "A Comparative Analysis of Metabarcoding and Morphology-based Identification of Benthic Communities across Different Regional Seas." *Ecology and Evolution* 8 (17): 8908–20.
<https://doi.org/10.1002/ece3.4283>.
- Carraro, Luca, Julian B. Stauffer, and Florian Altermatt. 2021. "How to Design Optimal eDNA Sampling Strategies for Biomonitoring in River Networks." *Environmental DNA* 3 (1): 157–72. <https://doi.org/10.1002/edn3.137>.
- Carvalho, Carolina S., Marina Elisa de Oliveira, Karen Giselle Rodriguez-Castro, Bruno H. Saranholi, and Pedro M. Galetti. 2021. "Efficiency of eDNA and iDNA in Assessing Vertebrate Diversity and Its Abundance." *Molecular Ecology Resources*, November, 1755-0998.13543. <https://doi.org/10.1111/1755-0998.13543>.
- Chang, Jia Jin Marc, Yin Cheong Aden Ip, Chin Soon Lionel Ng, and Danwei Huang. 2020. "Takeaways from Mobile DNA Barcoding with BentoLab and MinION." *Genes* 11 (10): 1121. <https://doi.org/10.3390/genes11101121>.

- Chua, Physilia Y.S., Sarah J. Bourlat, Cameron Ferguson, Petra Korlevic, Leia Zhao, Torbjørn Ekrem, Rudolf Meier, and Mara K.N. Lawniczak. 2023. "Future of DNA-Based Insect Monitoring." *Trends in Genetics* 39 (7): 531–44. <https://doi.org/10.1016/j.tig.2023.02.012>.
- Closek, Collin J., Jarrod A. Santora, Hilary A. Starks, Isaac D. Schroeder, Elizabeth A. Andruszkiewicz, Keith M. Sakuma, Steven J. Bograd, Elliott L. Hazen, John C. Field, and Alexandria B. Boehm. 2019. "Marine Vertebrate Biodiversity and Distribution Within the Central California Current Using Environmental DNA (eDNA) Metabarcoding and Ecosystem Surveys." *Frontiers in Marine Science* 6 (December): 732. <https://doi.org/10.3389/fmars.2019.00732>.
- Colas, F., M. Tardivel, J. Perchoc, M. Lunven, B. Forest, G. Guyader, M.M. Danielou, et al. 2018. "The ZooCAM, a New in-Flow Imaging System for Fast Onboard Counting, Sizing and Classification of Fish Eggs and Metazooplankton." *Progress in Oceanography* 166 (September): 54–65. <https://doi.org/10.1016/j.pocean.2017.10.014>.
- Cooper, Keith M, and Hubert L Rees. 2002. "National Marine Biological Analytical Quality Control Scheme (NMBAQC) Review of Standard Operating Procedures." 13. Science Series Aquatic Environmental Protection: Analytical Methods. CEFAS.
- Cordier, Tristan. 2019. "Bacterial Communities' Taxonomic and Functional Turnovers Both Accurately Predict Marine Benthic Ecological Quality Status." *Environmental DNA* 2 (2): 175–83. <https://doi.org/10.1002/edn3.55>.
- Cordier, Tristan, Laura Alonso-Sáez, Laure Apothéloz-Perret-Gentil, Eva Aylagas, David A. Bohan, Agnès Bouchez, Anthony Chariton, et al. 2020. "Ecosystems Monitoring Powered by Environmental Genomics: A Review of Current Strategies with an Implementation Roadmap." *Molecular Ecology*, June, mec.15472. <https://doi.org/10.1111/mec.15472>.
- Cordier, Tristan, Philippe Esling, Franck Lejzerowicz, Joana Visco, Amine Ouadahi, Catarina Martins, Tomas Cedhagen, and Jan Pawlowski. 2017. "Predicting the Ecological Quality Status of Marine Environments from eDNA Metabarcoding Data Using Supervised Machine Learning." *Environmental Science & Technology* 51 (16): 9118–26. <https://doi.org/10.1021/acs.est.7b01518>.
- Cordier, Tristan, Dominik Forster, Yoann Dufresne, Catarina I. M. Martins, Thorsten Stoeck, and Jan Pawlowski. 2018. "Supervised Machine Learning Outperforms Taxonomy-Based Environmental DNA Metabarcoding Applied to Biomonitoring." *Molecular Ecology Resources* 18 (6): 1381–91. <https://doi.org/10.1111/1755-0998.12926>.
- Cowart, Dominique A., Miguel Pinheiro, Olivier Mouchel, Marion Maguer, Jacques Grall, Jacques Miné, and Sophie Arnaud-Haond. 2015. "Metabarcoding Is Powerful yet Still Blind: A Comparative Analysis of Morphological and Molecular Surveys of Seagrass Communities." Edited by Silvia Mazzuca. *PLOS ONE* 10 (2): e0117562. <https://doi.org/10.1371/journal.pone.0117562>.
- Cristescu, Melania E. 2019. "Can Environmental RNA Revolutionize Biodiversity Science?" *Trends in Ecology & Evolution* 34: 694–97.
- Danielsen, Jørgen Skyrud, Luis Morgado, Sunil Mundra, Line Nybakken, Marie Davey, and Håvard Kauserud. 2021. "Establishment of Spruce Plantations in Native Birch Forests Reduces Soil Fungal Diversity." *FEMS Microbiology Ecology*.
- Davy, Christina M., Anne G. Kidd, and Chris C. Wilson. 2015. "Development and Validation of Environmental DNA (eDNA) Markers for Detection of Freshwater Turtles." Edited by Andrew R. Mahon. *PLOS ONE* 10 (7): e0130965. <https://doi.org/10.1371/journal.pone.0130965>.
- Deagle, Bruce E., Laurence J. Clarke, John A. Kitchener, Andrea M. Polanowski, and Andrew T. Davidson. 2018. "Genetic Monitoring of Open Ocean Biodiversity: An

- Evaluation of DNA Metabarcoding for Processing Continuous Plankton Recorder Samples." *Molecular Ecology Resources* 18 (3): 391–406. <https://doi.org/10.1111/1755-0998.12740>.
- Deiner, Kristy, Holly M. Bik, Elvira Mächler, Mathew Seymour, Anaïs Lacoursière-Roussel, Florian Altermatt, Simon Creer, et al. 2017. "Environmental DNA Metabarcoding: Transforming How We Survey Animal and Plant Communities." *Molecular Ecology* 26 (21): 5872–95. <https://doi.org/10.1111/mec.14350>.
- Di Muri, Cristina, Lori Lawson Handley, Colin W. Bean, Marco Benucci, Lynsey R. Harper, Ben James, Jianlong Li, Ian J. Winfield, and Bernd Hänfling. 2022. "Spatio-temporal Monitoring of Lake Fish Spawning Activity Using Environmental DNA Metabarcoding." *Environmental DNA*, July, edn3.343. <https://doi.org/10.1002/edn3.343>.
- Di Muri, Cristina, Lori Lawson Handley, Colin W. Bean, Jianlong Li, Graeme Peirson, Graham S. Sellers, Kerry Walsh, Hayley V. Watson, Ian J. Winfield, and Bernd Hänfling. 2020. "Read Counts from Environmental DNA (eDNA) Metabarcoding Reflect Fish Abundance and Biomass in Drained Ponds." *Metabarcoding and Metagenomics* 4 (October): e56959. <https://doi.org/10.3897/mbmg.4.56959>.
- DiBattista, Joseph D., James D. Reimer, Michael Stat, Giovanni D. Masucci, Piera Biondi, Maarten De Brauwer, Shaun P. Wilkinson, Anthony A. Chariton, and Michael Bunce. 2020. "Environmental DNA Can Act as a Biodiversity Barometer of Anthropogenic Pressures in Coastal Ecosystems." *Scientific Reports* 10 (1): 8365. <https://doi.org/10.1038/s41598-020-64858-9>.
- Djurhuus, Anni, Kathleen Pitz, Natalie A. Sawaya, Jaimie Rojas-Márquez, Brianna Michaud, Enrique Montes, Frank Muller-Karger, and Mya Breitbart. 2018. "Evaluation of Marine Zooplankton Community Structure through Environmental DNA Metabarcoding." *Limnology and Oceanography: Methods* 16 (4): 209–21. <https://doi.org/10.1002/lom3.10237>.
- Doble, Christopher J., Helen Hipperson, Walter Salzburger, Gavin J. Horsburgh, Chacha Mwita, David J. Murrell, and Julia J. Day. 2020. "Testing the Performance of Environmental DNA Metabarcoding for Surveying Highly Diverse Tropical Fish Communities: A Case Study from Lake Tanganyika." *Environmental DNA* 2 (1): 24–41. <https://doi.org/10.1002/edn3.43>.
- Doi, Hideyuki, Yoshihisa Akamatsu, Yutaka Watanabe, Masuji Goto, Ryutei Inui, Izumi Katano, Mariko Nagano, Teruhiko Takahara, and Toshifumi Minamoto. 2017. "Water Sampling for Environmental DNA Surveys by Using an Unmanned Aerial Vehicle: Drone Water Sampling for eDNA." *Limnology and Oceanography: Methods* 15 (11): 939–44. <https://doi.org/10.1002/lom3.10214>.
- Dougherty, Matthew M., Eric R. Larson, Mark A. Renshaw, Crysta A. Gantz, Scott P. Egan, Daniel M. Erickson, and David M. Lodge. 2016. "Environmental DNA (eDNA) Detects the Invasive Rusty Crayfish *Orconectes Rusticus* at Low Abundances." Edited by Chris Frid. *Journal of Applied Ecology* 53 (3): 722–32. <https://doi.org/10.1111/1365-2664.12621>.
- Drummond, Alexei J., Richard D. Newcomb, Thomas R. Buckley, Dong Xie, Andrew Dopheide, Benjamin CM Potter, Joseph Heled, et al. 2015. "Evaluating a Multigene Environmental DNA Approach for Biodiversity Assessment." *GigaScience* 4 (1): 46. <https://doi.org/10.1186/s13742-015-0086-1>.
- Duarte, Sofia, Barbara Leite, Maria Feio, Filipe Costa, and Ana Filipe. 2021. "Integration of DNA-Based Approaches in Aquatic Ecological Assessment Using Benthic Macroinvertebrates." *Water* 13 (3): 331. <https://doi.org/10.3390/w13030331>.
- Dubinsky, Eric A., Mark E. Conrad, Romy Chakraborty, Markus Bill, Sharon E. Borglin, James T. Hollibaugh, Olivia U. Mason, et al. 2013. "Succession of Hydrocarbon-

- Degrading Bacteria in the Aftermath of the *Deepwater Horizon* Oil Spill in the Gulf of Mexico.” *Environmental Science & Technology* 47 (19): 10860–67. <https://doi.org/10.1021/es401676y>.
- Dutra, Joyce, Glen García, Rosimeire Gomes, Mariana Cardoso, Árley Côrtes, Tales Silva, Luís De Jesus, et al. 2023. “Effective Biocorrosive Control in Oil Industry Facilities: 16S rRNA Gene Metabarcoding for Monitoring Microbial Communities in Produced Water.” *Microorganisms* 11 (4): 846. <https://doi.org/10.3390/microorganisms11040846>.
- Egeter, Bastian, Sara Peixoto, José C. Brito, Simon Jarman, Pamela Puppo, and Guillermo Velo-Antón. 2018. “Challenges for Assessing Vertebrate Diversity in Turbid Saharan Water-Bodies Using Environmental DNA.” Edited by Xavier Pochon. *Genome* 61 (11): 807–14. <https://doi.org/10.1139/gen-2018-0071>.
- Elbrecht, Vasco, and Florian Leese. 2017. “Validation and Development of COI Metabarcoding Primers for Freshwater Macroinvertebrate Bioassessment.” *Frontiers in Environmental Science* 5 (April). <https://doi.org/10.3389/fenvs.2017.00011>.
- Elbrecht, Vasco, Ecaterina Edith Vamos, Kristian Meissner, Jukka Aroviita, and Florian Leese. 2017. “Assessing Strengths and Weaknesses of DNA Metabarcoding-based Macroinvertebrate Identification for Routine Stream Monitoring.” Edited by Douglas Yu. *Methods in Ecology and Evolution* 8 (10): 1265–75. <https://doi.org/10.1111/2041-210X.12789>.
- Ely, Taylor, Paul H. Barber, Lauren Man, and Zachary Gold. 2021. “Short-Lived Detection of an Introduced Vertebrate eDNA Signal in a Nearshore Rocky Reef Environment.” Edited by Hideyuki Doi. *PLOS ONE* 16 (6): e0245314. <https://doi.org/10.1371/journal.pone.0245314>.
- Environment Agency. 2021. “Understanding Ecosystems and Resilience Using DNA.” Bristol.
- European Parliament and Council. 2000. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy. Official Journal of the European Communities, L327, 1–73. <http://data.europa.eu/eli/dir/2000/60/oj>
- Everett, Meredith V., and Linda K. Park. 2018. “Exploring Deep-Water Coral Communities Using Environmental DNA.” *Deep-Sea Research Part II: Topical Studies in Oceanography* 150 (September 2017): 229–41. <https://doi.org/10.1016/j.dsr2.2017.09.008>.
- Fan, Juntao, Shuping Wang, Hong Li, Zhenguang Yan, Yizhang Zhang, Xin Zheng, and Pengyuan Wang. 2020. “Modeling the Ecological Status Response of Rivers to Multiple Stressors Using Machine Learning: A Comparison of Environmental DNA Metabarcoding and Morphological Data.” *Water Research* 183 (September): 116004. <https://doi.org/10.1016/j.watres.2020.116004>.
- Farrell, Jessica A, Liam Whitmore, Narges Mashkour, Devon R Rollinson Ramia, Rachel S Thomas, Catherine B Eastman, Brooke Burkhalter, et al. 2022. “Detection and Population Genomics of Sea Turtle Species via Non-invasive Environmental DNA Analysis of Nesting Beach Sand Tracks and Oceanic Water.” *Molecular Ecology Resources*, April, 1755-0998.13617. <https://doi.org/10.1111/1755-0998.13617>.
- Fast, Kayla, Anakela Popp, Patrick O Neil, Stuart McGregor, and Michael Sandel. 2020. “Surveillance of a Federally Protected Freshwater Fish Using Loop-Mediated Isothermal Amplification (LAMP) and eDNA.” Preprint. Preprints. <https://doi.org/10.22541/au.159586067.74431972>.
- Fediajevaite, Julija, Victoria Priestley, Richard Arnold, and Vincent Savolainen. 2021. “Meta-analysis Shows That Environmental DNA Outperforms Traditional Surveys, but

- Warrants Better Reporting Standards.” *Ecology and Evolution* 11 (9): 4803–15. <https://doi.org/10.1002/ece3.7382>.
- Feio, Maria João, Sónia R.Q. Serra, Andreia Mortágua, Agnès Bouchez, Frédéric Rimet, Valentin Vasselon, and Salomé F.P. Almeida. 2020. “A Taxonomy-Free Approach Based on Machine Learning to Assess the Quality of Rivers with Diatoms.” *Science of The Total Environment* 722 (June): 137900. <https://doi.org/10.1016/j.scitotenv.2020.137900>.
- Fernández, Sara, Saúl Rodríguez-Martínez, Jose L. Martínez, Eva Garcia-Vazquez, and Alba Ardura. 2019. “How Can eDNA Contribute in Riverine Macroinvertebrate Assessment? A Metabarcoding Approach in the Nalón River (Asturias, Northern Spain).” *Environmental DNA* 1 (4): 385–401. <https://doi.org/10.1002/edn3.40>.
- Ferreira, Diogo F, Crinan Jarrett, Patrick Jules Atagana, Luke L Powell, and Hugo Rebelo. 2021. “Are Bat Mist Nets Ideal for Capturing Bats? From Ultrathin to Bird Nets, a Field Test.” Edited by Amy Baird. *Journal of Mammalogy* 102 (6): 1627–34. <https://doi.org/10.1093/jmammal/gyab109>.
- Formel, Nathan, Ian C. Enochs, Chris Sinigalliano, Sean R. Anderson, and Luke R. Thompson. 2021. “Subsurface Automated Samplers for eDNA (SASe) for Biological Monitoring and Research.” *HardwareX* 10 (October): e00239. <https://doi.org/10.1016/j.ohx.2021.e00239>.
- Franklin, Thomas W., Kevin S. McKelvey, Jessie D. Golding, Daniel H. Mason, Joseph C. Dysthe, Kristine L. Pilgrim, John R. Squires, et al. 2019. “Using Environmental DNA Methods to Improve Winter Surveys for Rare Carnivores: DNA from Snow and Improved Noninvasive Techniques.” *Biological Conservation* 229 (January): 50–58. <https://doi.org/10.1016/j.biocon.2018.11.006>.
- Franz, Daniel. 2020. “Black and Yellow Letter Law: Managing Rusty Patched Bumble Bee Conservation Under the Endangered Species Act.” *ENVIRONMENTAL LAW REVIEW* 31.
- Frontalini, Fabrizio, Tristan Cordier, Eszter Balassi, Eric Armynot du Chatelet, Kristina Cermakova, Laure Apothéoz-Perret-Gentil, Maria Virginia Alves Martins, et al. 2020. “Benthic Foraminiferal Metabarcoding and Morphology-Based Assessment around Three Offshore Gas Platforms: Congruence and Complementarity.” *Environment International* 144 (November): 106049. <https://doi.org/10.1016/j.envint.2020.106049>.
- Frostegård, Åsa, Anders Tunlid, and Erland Bååth. 2011. “Use and Misuse of PLFA Measurements in Soils.” *Soil Biology and Biochemistry* 43 (8): 1621–25. <https://doi.org/10.1016/j.soilbio.2010.11.021>.
- Frühe, Larissa, Tristan Cordier, Verena Dully, Hans-Werner Breiner, Guillaume Lentendu, Jan Pawlowski, Catarina Martins, Thomas A. Wilding, and Thorsten Stoeck. 2020. “Supervised Machine Learning Is Superior to Indicator Value Inference in Monitoring the Environmental Impacts of Salmon Aquaculture Using eDNA Metabarcodes.” *Molecular Ecology* 30 (13): 2988–3006. <https://doi.org/10.1111/mec.15434>.
- Gogarten, Jan F., Constanze Hoffmann, Mimi Arandjelovic, Andreas Sachse, Kevin Merkel, Paula Dieguez, Anthony Agbor, et al. 2020. “Fly-derived DNA and Camera Traps Are Complementary Tools for Assessing Mammalian Biodiversity.” *Environmental DNA* 2 (1): 63–76. <https://doi.org/10.1002/edn3.46>.
- Gold, Zachary, Andrew Olaf Shelton, Helen R. Casendino, Joe Duprey, Ramón Gallego, Amy Van Cise, Mary Fisher, et al. 2023. “Signal and Noise in Metabarcoding Data.” Edited by Ruslan Kalendar. *PLOS ONE* 18 (5): e0285674. <https://doi.org/10.1371/journal.pone.0285674>.
- Golpour, Amin, Marek Smejkal, Cech Martin, Romulo dos Santos, Allan T Souza, Tomas Juza, Carlos Martinez, et al. 2022. “Similarities and Differences in Fish Community Composition Accessed by Electrofishing, Gill Netting, Seining, Trawling, and Water

- eDNA Metabarcoding in Temperate Reservoirs." *Frontiers in Ecology and Evolution* 10: 17.
- Gonzalo, Martin, and Maria de los Reyes Fernandez. 2012. "Diatoms as Indicators of Water Quality and Ecological Status: Sampling, Analysis and Some Ecological Remarks." In *Ecological Water Quality - Water Treatment and Reuse*, 183–204.
- Green, Hyatt C., Richard A. Haugland, Manju Varma, Hana T. Millen, Mark A. Borchardt, Katharine G. Field, William A. Walters, et al. 2014. "Improved HF183 Quantitative Real-Time PCR Assay for Characterization of Human Fecal Pollution in Ambient Surface Water Samples." Edited by F. E. Löffler. *Applied and Environmental Microbiology* 80 (10): 3086–94. <https://doi.org/10.1128/AEM.04137-13>.
- Guillerault, N., S. Bouletreau, A. Iribar, A. Valentini, and F. Santoul. 2017. "Application of DNA Metabarcoding on Faeces to Identify European Catfish *Silurus Glanis* Diet: Dna Metabarcoding of *s. Glanis* Faeces." *Journal of Fish Biology* 90 (5): 2214–19. <https://doi.org/10.1111/jfb.13294>.
- Günther, Babett, Thomas Knebelberger, Hermann Neumann, Silke Laakmann, and Pedro Martínez Arbizu. 2018. "Metabarcoding of Marine Environmental DNA Based on Mitochondrial and Nuclear Genes." *Scientific Reports* 8 (1): 14822. <https://doi.org/10.1038/s41598-018-32917-x>.
- Hallam, Jane, Elizabeth L. Clare, John Iwan Jones, and Julia J. Day. 2021. "Biodiversity Assessment across a Dynamic Riverine System: A Comparison of eDNA Metabarcoding versus Traditional Fish Surveying Methods." *Environmental DNA* 3 (6): 1247–66. <https://doi.org/10.1002/edn3.241>.
- Hänfling, Bernd, Lori Lawson Handley, Daniel S. Read, Christoph Hahn, Jianlong Li, Paul Nichols, Rosetta C. Blackman, Anna Oliver, and Ian J. Winfield. 2016. "Environmental DNA Metabarcoding of Lake Fish Communities Reflects Long-Term Data from Established Survey Methods." *Molecular Ecology* 25 (13): 3101–19. <https://doi.org/10.1111/mec.13660>.
- Hansen, Brian K., Magnus W. Jacobsen, Anne Lise Middelboe, Christina M. Preston, Roman Marin, Dorte Bekkevold, Steen W. Knudsen, Peter R. Møller, and Einar E. Nielsen. 2020. "Remote, Autonomous Real-Time Monitoring of Environmental DNA from Commercial Fish." *Scientific Reports* 10 (1): 13272. <https://doi.org/10.1038/s41598-020-70206-8>.
- Harper, Lynsey R., Lori Lawson Handley, Angus I. Carpenter, Muhammad Ghazali, Cristina Di Muri, Callum J. Macgregor, Thomas W. Logan, et al. 2019. "Environmental DNA (eDNA) Metabarcoding of Pond Water as a Tool to Survey Conservation and Management Priority Mammals." *Biological Conservation* 238 (October): 108225. <https://doi.org/10.1016/j.biocon.2019.108225>.
- Harper, Lynsey R., Lori Lawson Handley, Carl D. Sayer, Daniel S. Read, Marco Benucci, Rosetta C. Blackman, Matthew J. Hill, and Bernd Hänfling. 2021a. "Assessing the Impact of the Threatened Crucian Carp (*Carassius Carassius*) on Pond Invertebrate Diversity: A Comparison of Conventional and Molecular Tools." *Molecular Ecology* 30 (13): 3252–69. <https://doi.org/10.1111/mec.15670>.
- . 2021b. "Assessing the Impact of the Threatened Crucian Carp (*Carassius Carassius*) on Pond Invertebrate Diversity: A Comparison of Conventional and Molecular Tools." *Molecular Ecology* 30 (13): 3252–69. <https://doi.org/10.1111/mec.15670>.
- Hayami, Kana, Masayuki K. Sakata, Takashi Inagawa, Jiro Okitsu, Izumi Katano, Hideyuki Doi, Katsuki Nakai, et al. 2020. "Effects of Sampling Seasons and Locations on Fish Environmental DNA Metabarcoding in Dam Reservoirs." *Ecology and Evolution* 10 (12): 5354–67. <https://doi.org/10.1002/ece3.6279>.

- Hering, Daniel, Angel Borja, J.Iwan Jones, Didier Pont, Pieter Boets, Agnes Bouchez, Kat Bruce, et al. 2018. "Implementation Options for DNA-Based Identification into Ecological Status Assessment under the European Water Framework Directive." *Water Research* 138 (July): 192–205. <https://doi.org/10.1016/j.watres.2018.03.003>.
- Hervé, Alix, Isabelle Domaizon, Jean-Marc Baudoin, Tony Dejean, Pierre Gibert, Pauline Jean, Tiphaine Peroux, et al. 2022. "Spatio-Temporal Variability of eDNA Signal and Its Implication for Fish Monitoring in Lakes." Edited by Hideyuki Doi. *PLOS ONE* 17 (8): e0272660. <https://doi.org/10.1371/journal.pone.0272660>.
- Heyde, Mieke, Michael Bunce, Grant Wardell-Johnson, Kristen Fernandes, Nicole E. White, and Paul Nevill. 2020. "Testing Multiple Substrates for Terrestrial Biodiversity Monitoring Using Environmental DNA Metabarcoding." *Molecular Ecology Resources* 20 (3): 732–45. <https://doi.org/10.1111/1755-0998.13148>.
- Heyde, Mieke van der, Michael Bunce, and Paul Nevill. 2022. "Key Factors to Consider in the Use of Environmental DNA Metabarcoding to Monitor Terrestrial Ecological Restoration." *Science of The Total Environment* 848 (November): 157617. <https://doi.org/10.1016/j.scitotenv.2022.157617>.
- Hinz, Shawn, Jennifer Coston-Guarini, Michael Marnane, and Jean-Marc Guarini. 2022. "Evaluating eDNA for Use within Marine Environmental Impact Assessments," 27.
- Höss, S., E. Claus, P.C. Von Der Ohe, M. Brinke, H. Güde, P. Heininger, and W. Traunspurger. 2011. "Nematode Species at Risk — A Metric to Assess Pollution in Soft Sediments of Freshwaters." *Environment International* 37 (5): 940–49. <https://doi.org/10.1016/j.envint.2011.03.013>.
- Hunter, Margaret E., Sara J. Oyler-McCance, Robert M. Dorazio, Jennifer A. Fike, Brian J. Smith, Charles T. Hunter, Robert N. Reed, and Kristen M. Hart. 2015. "Environmental DNA (eDNA) Sampling Improves Occurrence and Detection Estimates of Invasive Burmese Pythons." Edited by Andrew R. Mahon. *PLOS ONE* 10 (4): e0121655. <https://doi.org/10.1371/journal.pone.0121655>.
- Ibarbalz, Federico M., Nicolas Henry, Manoela C. Brandão, Séverine Martini, Greta Busseni, Hannah Byrne, Luis Pedro Coelho, et al. 2019. "Global Trends in Marine Plankton Diversity across Kingdoms of Life." *Cell* 179 (5): 1084-1097.e21. <https://doi.org/10.1016/j.cell.2019.10.008>.
- IOGP. 2020. "Environmental Sampling and Monitoring from Airborne and Satellite Remote Sensing," 68.
- Ip, Yin Cheong Aden, Jia Jin Marc Chang, Ren Min Oh, Zheng Bin Randolph Quek, Yong Kit Samuel Chan, Andrew G. Bauman, and Danwei Huang. 2023. "Seq' and ARMS Shall Find: DNA (Meta)Barcoding of Autonomous Reef Monitoring Structures across the Tree of Life Uncovers Hidden Cryptobiome of Tropical Urban Coral Reefs." *Molecular Ecology* 32 (23): 6223–42. <https://doi.org/10.1111/mec.16568>.
- Irfan Ali Phulpoto. 2021. "Effect of Natural Microbiome and Culturable Biosurfactants-Producing Bacterial Consortia of Freshwater Lake on Petroleum-Hydrocarbon Degradation." Translated by Irfan Ali Phulpoto, Bowen Hu, Yanfen Wang, Fabrice Ndayisenga, Jinmei Li, and Zhisheng Yu. *Science of the Total Environment* v. 751: 141720-. <https://doi.org/10.1016/j.scitotenv.2020.141720>.
- Jac, Cyrielle, Nicolas Desroy, Jean-Claude Duchêne, Aurélie Foveau, Céline Labrune, Lyvia Lescure, and Sandrine Vaz. 2021. "Assessing the Impact of Trawling on Benthic Megafauna: Comparative Study of Video Surveys vs. Scientific Trawling." Edited by Joanna Norkko. *ICES Journal of Marine Science* 78 (5): 1636–49. <https://doi.org/10.1093/icesjms/fsab033>.
- Jamy, Mahwash, Rachel Foster, Pierre Barbera, Lucas Czech, Alexey Kozlov, Alexandros Stamatakis, David Bass, and Fabien Burki. 2020. "Long Metabarcoding of the Eukaryotic rDNA Operon to Phylogenetically and Taxonomically Resolve

- Environmental Diversity.” *Molecular Ecology Resources* 20: 429–43.
<https://doi.org/10.1101/627828>.
- Janssen, Peter H., Penelope S. Yates, Bronwyn E. Grinton, Paul M. Taylor, and Michelle Sait. 2002. “Improved Culturability of Soil Bacteria and Isolation in Pure Culture of Novel Members of the Divisions *Acidobacteria*, *Actinobacteria*, *Proteobacteria*, and *Verrucomicrobia*.” *Applied and Environmental Microbiology* 68 (5): 2391–96.
<https://doi.org/10.1128/AEM.68.5.2391-2396.2002>.
- Jeunen, Gert-Jan, Ulla von Ammon, Hugh Cross, Sara Ferreira, Miles Lamare, Robert Day, Jackson Treece, et al. 2022. “Moving Environmental DNA (eDNA) Technologies from Benchtop to the Field Using Passive Sampling and PDQEX Extraction.” *Environmental DNA* 4 (6): 1420–33. <https://doi.org/10.1002/edn3.356>.
- Jeunen, Gert-Jan, Miles D. Lamare, Michael Knapp, Hamish G. Spencer, Helen R. Taylor, Michael Stat, Michael Bunce, and Neil J. Gemmill. 2020. “Water Stratification in the Marine Biome Restricts Vertical Environmental DNA (eDNA) Signal Dispersal.” *Environmental DNA* 2 (1): 99–111. <https://doi.org/10.1002/edn3.49>.
- Jin, Decai, Xiao Kong, Bingjian Cui, Shulan Jin, Yunfeng Xie, Xingrun Wang, and Ye Deng. 2018. “Bacterial Communities and Potential Waterborne Pathogens within the Typical Urban Surface Waters.” *Scientific Reports* 8 (1): 13368.
<https://doi.org/10.1038/s41598-018-31706-w>.
- JNCC. 2017. “JNCC Guidelines for Minimising the Risk of Injury to Marine Mammals from Geophysical Surveys.”
- Johnson, Mark D., Matthew A. Barnes, Nina R. Garrett, and Elizabeth L. Clare. 2023. “Answers Blowing in the Wind: Detection of Birds, Mammals, and Amphibians with Airborne Environmental DNA in a Natural Environment over a Yearlong Survey.” *Environmental DNA* 5 (2): 375–87. <https://doi.org/10.1002/edn3.388>.
- Johnson, Mark D., Mohamed Fokar, Robert D. Cox, and Matthew A. Barnes. 2021. “Airborne Environmental DNA Metabarcoding Detects More Diversity, with Less Sampling Effort, than a Traditional Plant Community Survey.” *BMC Ecology and Evolution* 21 (1): 218. <https://doi.org/10.1186/s12862-021-01947-x>.
- Johnson, Wilson H., Marlis R. Douglas, Jeffrey A. Lewis, Tara N. Stuecker, Franck G. Carbonero, Bradley J. Austin, Michelle A. Evans-White, Sally A. Entekin, and Michael E. Douglas. 2017. “Do Biofilm Communities Respond to the Chemical Signatures of Fracking? A Test Involving Streams in North-Central Arkansas.” *BMC Microbiology* 17 (1): 29. <https://doi.org/10.1186/s12866-017-0926-5>.
- Jungbluth, Michelle J., Katharine M. Hanson, Petra H. Lenz, H. Eve Robinson, and Erica Goetze. 2022. “Species-specific Biomass Estimation from Gene Copy Number in Metazoan Plankton.” *Limnology and Oceanography: Methods* 20 (6): 305–19.
<https://doi.org/10.1002/lom3.10487>.
- Jurelevicius, Diogo, Vanessa Marques Alvarez, Joana Montezano Marques, Laryssa Ribeiro Fonseca de Sousa Lima, Felipe de Almeida Dias, and Lucy Seldin. 2013. “Bacterial Community Response to Petroleum Hydrocarbon Amendments in Freshwater, Marine, and Hypersaline Water-Containing Microcosms.” *Applied and Environmental Microbiology* 79 (19): 5927–35. <https://doi.org/10.1128/AEM.02251-13>.
- Keck, François, Rosetta C. Blackman, Raphael Bossart, Jeanine Brantschen, Marjorie Couton, Samuel Hürlemann, Dominik Kirschner, Nadine Locher, Heng Zhang, and Florian Altermatt. 2022. “Meta-Analysis Shows Both Congruence and Complementarity of DNA Metabarcoding to Traditional Methods for Biological Community Assessment.” *Molecular Ecology* 31 (6).
<https://doi.org/10.1101/2021.06.29.450286>.
- Keck, François, Samuel Hürlemann, Nadine Locher, Christian Stamm, Kristy Deiner, and Florian Altermatt. 2022. “A Triad of Kicknet Sampling, eDNA Metabarcoding, and

- Predictive Modeling to Assess Richness of Mayflies, Stoneflies and Caddisflies in Rivers.” *Metabarcoding and Metagenomics* 6 (May): e79351.
<https://doi.org/10.3897/mbmg.6.79351>.
- Keeley, Nigel, Olivier Laroche, Murray Birch, and Xavier Pochon. 2021. “A Substrate-Independent Benthic Sampler (SIBS) for Hard and Mixed-Bottom Marine Habitats: A Proof-of-Concept Study.” *Frontiers in Marine Science* 8 (April): 627687.
<https://doi.org/10.3389/fmars.2021.627687>.
- Keeley, Nigel, Susanna A. Wood, and Xavier Pochon. 2018. “Development and Preliminary Validation of a Multi-Trophic Metabarcoding Biotic Index for Monitoring Benthic Organic Enrichment.” *Ecological Indicators* 85 (February): 1044–57.
<https://doi.org/10.1016/j.ecolind.2017.11.014>.
- Kelly, M. G., A. Cazaubon, E. Coring, A. Dell’Uomo, L. Ector, B. Goldsmith, H. Guasch, et al. 1998. “Recommendations for the Routine Sampling of Diatoms for Water Quality Assessments in Europe.” *Journal of Applied Phycology* 10 (2): 215–24.
<https://doi.org/10.1023/A:1008033201227>.
- Kelly, Martyn, Neil Boonham, Steve Juggins, David Mann, and Rachel Glover. 2020. “Further Development of a DNA Based Metabarcoding Approach to Assess Diatom Communities in Rivers.” SC160014/R. Chief Scientist’s Group Report. Environment Agency.
https://assets.publishing.service.gov.uk/media/61a0def28fa8f5037778c1f4/Assessing_river_nutrients_using_diatom_DNA_-_further_development_of_an_operational_method_-_report.pdf.
- Kim, Park, Jo, and Kwak. 2019. “Comparison of Water Sampling between Environmental DNA Metabarcoding and Conventional Microscopic Identification: A Case Study in Gwangyang Bay, South Korea.” *Applied Sciences* 9 (16): 3272.
<https://doi.org/10.3390/app9163272>.
- Kirse, Ameli, Sarah J. Bourlat, Kathrin Langen, and Vera G. Fonseca. 2021. “Metabarcoding Malaise Traps and Soil eDNA Reveals Seasonal and Local Arthropod Diversity Shifts.” *Scientific Reports* 11 (1): 10498. <https://doi.org/10.1038/s41598-021-89950-6>.
- Kirtane, Anish, John D. Atkinson, and Lauren Sassoubre. 2020. “Design and Validation of Passive Environmental DNA Samplers Using Granular Activated Carbon and Montmorillonite Clay.” *Environmental Science & Technology* 54 (19): 11961–70.
<https://doi.org/10.1021/acs.est.0c01863>.
- Kitagami, Yudai, and Yosuke Matsuda. 2022. “High-Throughput Sequencing Covers Greater Nematode Diversity than Conventional Morphotyping on Natural Cedar Forests in Yakushima Island, Japan.” *SSRN Electronic Journal*.
<https://doi.org/10.2139/ssrn.4144031>.
- Klunder, Lise, Marc S S Lavaleye, Amalia Filippidi, and Gert-Jan Reichart. 2018. “Impact of an Artificial Structure on the Benthic Community Composition in the Southern North Sea: Assessed by a Morphological and Molecular Approach.” *ICES Journal of Marine Science*, 11.
- Klunder, Lise, Henko de Stigter, Marc S. S. Lavaleye, Judith D. L. van Bleijswijk, Henk W. van der Veer, Gert-Jan Reichart, and Gerard C. A. Duineveld. 2020. “A Molecular Approach to Explore the Background Benthic Fauna Around a Hydrothermal Vent and Their Larvae: Implications for Future Mining of Deep-Sea SMS Deposits.” *Frontiers in Marine Science* 7 (March): 134.
<https://doi.org/10.3389/fmars.2020.00134>.
- Klymus, Katy E., Catherine A. Richter, Nathan Thompson, and Jo Ellen Hinck. 2017. “Metabarcoding of Environmental DNA Samples to Explore the Use of Uranium Mine Containment Ponds as a Water Source for Wildlife.” *Diversity* 9 (4): 54.
<https://doi.org/10.3390/d9040054>.

- Knapik, Kamila, Andrea Bagi, Adriana Krolicka, and Thierry Baussant. 2020. "Metatranscriptomic Analysis of Oil-Exposed Seawater Bacterial Communities Archived by an Environmental Sample Processor (ESP)." *Microorganisms* 8 (5): 744. <https://doi.org/10.3390/microorganisms8050744>.
- Kowarski, Katie A., Briand J. Gaudet, Arthur J. Cole, Emily E Maxner, Stephen P Turner, S. Bruce Martin, Hansen D. Johnson, and John E. Moloney. 2020. "Near Real-Time Marine Mammal Monitoring from Gliders: Practical Challenges, System Development, and Management Implications." *The Journal of the Acoustical Society of America* 148 (3): 1215–30. <https://doi.org/10.1121/10.0001811>.
- Koziol, Adam, Michael Stat, Tiffany Simpson, Simon Jarman, Joseph D. DiBattista, Euan S. Harvey, Michael Marnane, Justin McDonald, and Michael Bunce. 2019. "Environmental DNA Metabarcoding Studies Are Critically Affected by Substrate Selection." *Molecular Ecology Resources* 19 (2): 366–76. <https://doi.org/10.1111/1755-0998.12971>.
- Kranzfelder, Petra, Alyssa M. Anderson, Alexander T. Egan, Jane E. Mazack, Jr. Bouchard R. William, Moriya M. Rufer, and Jr. Ferrington Leonard C. 2015. "Use of Chironomidae (Diptera) Surface-Floating Pupal Exuviae as a Rapid Bioassessment Protocol for Water Bodies." *JoVE*, no. 101 (July): e52558. <https://doi.org/10.3791/52558>.
- Krehenwinkel, Henrik, Aaron Pomerantz, James B Henderson, Susan R Kennedy, Jun Ying Lim, Varun Swamy, Juan Diego Shoobridge, et al. 2019. "Nanopore Sequencing of Long Ribosomal DNA Amplicons Enables Portable and Simple Biodiversity Assessments with High Phylogenetic Resolution across Broad Taxonomic Scale," no. January: 1–16. <https://doi.org/10.1093/gigascience/giz006>.
- Krolicka, Adriana, Catherine Boccadoro, Mari Mæland Nilsen, and Thierry Baussant. 2017. "Capturing Early Changes in the Marine Bacterial Community as a Result of Crude Oil Pollution in a Mesocosm Experiment." *Microbes and Environments* 32 (4): 358–66. <https://doi.org/10.1264/jsme2.ME17082>.
- Krolicka, Adriana, Catherine Boccadoro, Mari Mæland Nilsen, Elif Demir-Hilton, Jim Birch, Christina Preston, Chris Scholin, and Thierry Baussant. 2019. "Identification of Microbial Key-Indicators of Oil Contamination at Sea through Tracking of Oil Biotransformation: An Arctic Field and Laboratory Study." *Science of The Total Environment* 696 (December): 133715. <https://doi.org/10.1016/j.scitotenv.2019.133715>.
- Krolicka, Adriana, Catherine Boccadoro, Christina M. Preston, and Thierry Baussant. 2014. "Detection of Oil Leaks by Quantifying Hydrocarbonoclastic Bacteria in Cold Marine Environments Using the Environmental Sample Processor." In *Proceedings of the 37th AMOP Technical Seminar on Environmental Contamination and Response*.
- Krolicka, Adriana, Alessio Gomiero, and Thierry Baussant. 2020. "qPCR-Based Assessment of Microfaunal Indicators of Oil for Monitoring Benthos around Oil and Gas Platforms." *Science of The Total Environment* 736 (September): 139527. <https://doi.org/10.1016/j.scitotenv.2020.139527>.
- Kuwae, Michinobu, Hiromichi Tamai, Hideyuki Doi, Masayuki K. Sakata, Toshifumi Minamoto, and Yoshiaki Suzuki. 2020. "Sedimentary DNA Tracks Decadal-Centennial Changes in Fish Abundance." *Communications Biology* 3 (1): 558. <https://doi.org/10.1038/s42003-020-01282-9>.
- Kyle, Kathleen E., Michael C. Allen, Jeffrey Dragon, John F. Bunnell, Howard K. Reinert, Robert Zappalorti, Benjamin D. Jaffe, Jordan C. Angle, and Julie L. Lockwood. 2022. "Combining Surface and Soil Environmental DNA with Artificial Cover Objects to Improve Terrestrial Reptile Survey Detection." *Conservation Biology* 36 (6). <https://doi.org/10.1111/cobi.13939>.

- Lacoursière-Roussel, Anaïs, Yohann Dubois, Eric Normandeau, and Louis Bernatchez. 2016. "Improving Herpetological Surveys in Eastern North America Using the Environmental DNA Method." Edited by Sarah Adamowicz. *Genome* 59 (11): 991–1007. <https://doi.org/10.1139/gen-2015-0218>.
- Lafont, Michel, Guillaume Tixier, Jiri Marsalek, Céline Jézéquel, Pascal Breil, and Laurent Schmitt. 2012. "From Research to Operational Biomonitoring of Freshwaters: A Suggested Conceptual Framework and Practical Solutions." *Ecohydrology & Hydrobiology* 12 (1): 9–20. <https://doi.org/10.2478/v10104-012-0004-4>.
- Lamendella, Regina, Steven Strutt, Sharon Borglin, Romy Chakraborty, Neslihan Tas, Olivia U. Mason, Jenni Hultman, Emmanuel Prestat, Terry C. Hazen, and Janet K. Jansson. 2014. "Assessment of the Deepwater Horizon Oil Spill Impact on Gulf Coast Microbial Communities." *Frontiers in Microbiology* 5 (April). <https://doi.org/10.3389/fmicb.2014.00130>.
- Lamy, Thomas, Kathleen J. Pitz, Francisco P. Chavez, Christie E. Yorke, and Robert J. Miller. 2021. "Environmental DNA Reveals the Fine-Grained and Hierarchical Spatial Structure of Kelp Forest Fish Communities." *Scientific Reports* 11 (1): 14439. <https://doi.org/10.1038/s41598-021-93859-5>.
- Lanzén, Anders, Thomas G. Dahlgren, Andrea Bagi, and Jon Thomassen Hestetun. 2021. "Benthic eDNA Metabarcoding Provides Accurate Assessments of Impact from Oil Extraction, and Ecological Insights." *Ecological Indicators* 130 (November): 108064. <https://doi.org/10.1016/j.ecolind.2021.108064>.
- Lanzén, Anders, Katrine Lekang, Inge Jonassen, Eric M. Thompson, and Christofer Troedsson. 2016. "High-Throughput Metabarcoding of Eukaryotic Diversity for Environmental Monitoring of Offshore Oil-Drilling Activities." *Molecular Ecology* 25 (17): 4392–4406. <https://doi.org/10.1111/mec.13761>.
- Lanzén, Anders, Iñaki Mendibil, Ángel Borja, and Laura Alonso-Sáez. 2021. "A Microbial *Mandala* for Environmental Monitoring: Predicting Multiple Impacts on Estuarine Prokaryote Communities of the Bay of Biscay." *Molecular Ecology* 30 (13): 2969–87. <https://doi.org/10.1111/mec.15489>.
- Laroche, Olivier, Oliver Kersten, Craig R. Smith, and Erica Goetze. 2020a. "From Sea Surface to Seafloor: A Benthic Allochthonous eDNA Survey for the Abyssal Ocean," May. <https://doi.org/10.1101/2020.05.07.082602>.
- . 2020b. "Environmental DNA Surveys Detect Distinct Metazoan Communities across Abyssal Plains and Seamounts in the Western Clarion Clipperton Zone." *Molecular Ecology* 29 (23): 4588–4604. <https://doi.org/10.1111/mec.15484>.
- Laroche, Olivier, Susanna A. Wood, Louis A. Tremblay, Joanne I. Ellis, Gavin Lear, and Xavier Pochon. 2018. "A Cross-Taxa Study Using Environmental DNA/RNA Metabarcoding to Measure Biological Impacts of Offshore Oil and Gas Drilling and Production Operations." *Marine Pollution Bulletin* 127 (February): 97–107. <https://doi.org/10.1016/j.marpolbul.2017.11.042>.
- Laroche, Olivier, Susanna A. Wood, Louis A. Tremblay, Joanne I. Ellis, Franck Lejzerowicz, Jan Pawlowski, Gavin Lear, Javier Atalah, and Xavier Pochon. 2016. "First Evaluation of Foraminiferal Metabarcoding for Monitoring Environmental Impact from an Offshore Oil Drilling Site." *Marine Environmental Research* 120 (September): 225–35. <https://doi.org/10.1016/j.marenvres.2016.08.009>.
- Laroche, Olivier, Susanna A. Wood, Louis A. Tremblay, Gavin Lear, Joanne I. Ellis, and Xavier Pochon. 2017. "Metabarcoding Monitoring Analysis: The Pros and Cons of Using Co-Extracted Environmental DNA and RNA Data to Assess Offshore Oil Production Impacts on Benthic Communities." *PeerJ* 5 (May): e3347. <https://doi.org/10.7717/peerj.3347>.

- Lawson Handley, Lori, Daniel S. Read, Ian J. Winfield, Helen Kimbell, Harriet Johnson, Jianlong Li, Christoph Hahn, et al. 2019. "Temporal and Spatial Variation in Distribution of Fish Environmental DNA in England's Largest Lake." *Environmental DNA* 1 (1): 26–39. <https://doi.org/10.1002/edn3.5>.
- Le, Jennifer T., Lisa A. Levin, Franck Lejzerowicz, Tristan Cordier, Andrew J. Gooday, and Jan Pawlowski. 2022. "Scientific and Budgetary Trade-offs between Morphological and Molecular Methods for Deep-sea Biodiversity Assessment." *Integrated Environmental Assessment and Management* 18 (3): 655–63. <https://doi.org/10.1002/ieam.4466>.
- Leempoel, Kevin, Trevor Hebert, and Elizabeth A. Hadly. 2020. "A Comparison of eDNA to Camera Trapping for Assessment of Terrestrial Mammal Diversity." *Proceedings of the Royal Society B: Biological Sciences* 287 (1918). <https://doi.org/10.1098/rspb.2019.2353>.
- Lejzerowicz, Franck, Philippe Esling, Loïc Pillet, Thomas A. Wilding, Kenneth D. Black, and Jan Pawlowski. 2015. "High-Throughput Sequencing and Morphology Perform Equally Well for Benthic Monitoring of Marine Ecosystems." *Scientific Reports* 5 (1): 13932. <https://doi.org/10.1038/srep13932>.
- Lekang, Katrine, Anders Lanzén, Inge Jonassen, Eric Thompson, and Christofer Troedsson. 2020. "Evaluation of a Eukaryote Phylogenetic Microarray for Environmental Monitoring of Marine Sediments." *Marine Pollution Bulletin* 154 (May): 111102. <https://doi.org/10.1016/j.marpolbul.2020.111102>.
- Li, Jianlong, Tristan W Hatton-Ellis, Lori Lawson Handley, Helen S Kimbell, Marco Benucci, Graeme Peirson, and Bernd Hänfling. 2019. "Ground-Truthing of a Fish-Based Environmental DNA Metabarcoding Method for Assessing the Quality of Lakes." *Journal of Applied Ecology* 56 (5).
- Li, Min, Ting Lei, Guobin Wang, Danli Zhang, Huaxi Liu, and Zhiwei Zhang. 2023. "Monitoring Insect Biodiversity and Comparison of Sampling Strategies Using Metabarcoding: A Case Study in the Yanshan Mountains, China." *Ecology and Evolution* 13 (4): e10031. <https://doi.org/10.1002/ece3.10031>.
- Littlefair, Joanne E., Lee E. Hrenchuk, Paul J. Blanchfield, Michael D. Rennie, and Melania E. Cristescu. 2020. "Thermal Stratification and Fish Thermal Preference Explain Vertical eDNA Distributions in Lakes." *Molecular Ecology* 30 (13): 3083–96. <https://doi.org/10.1111/mec.15623>.
- Liu, Zifang, Rupert A. Collins, Charles Baillie, Sophie Rainbird, Rachel Brittain, Andrew M. Griffiths, David W. Sims, Stefano Mariani, and Martin J. Genner. 2022. "Environmental DNA Captures Elasmobranch Diversity in a Temperate Marine Ecosystem." *Environmental DNA* 4 (5): 1024–38. <https://doi.org/10.1002/edn3.294>.
- Loos, Luna M., and Reindert Nijland. 2021. "Biases in Bulk: DNA Metabarcoding of Marine Communities and the Methodology Involved." *Molecular Ecology* 30 (13): 3270–88. <https://doi.org/10.1111/mec.15592>.
- Luo, Mingjie, Yinqiu Ji, David Warton, and Douglas W. Yu. 2022. "Extracting Abundance Information from DNA -based Data." *Molecular Ecology Resources*, August, 1755-0998.13703. <https://doi.org/10.1111/1755-0998.13703>.
- Lyet, Arnaud, Loïc Pellissier, Alice Valentini, Tony Dejean, Abigail Hehmeyer, and Robin Naidoo. 2021. "eDNA Sampled from Stream Networks Correlates with Camera Trap Detection Rates of Terrestrial Mammals." *Scientific Reports* 11 (1): 11362. <https://doi.org/10.1038/s41598-021-90598-5>.
- Lynggaard, Christina, Mads Frost Bertelsen, Casper V. Jensen, Matthew S. Johnson, Tobias Guldberg Frøslev, Morten Tange Olsen, and Kristine Bohmann. 2022. "Airborne Environmental DNA for Terrestrial Vertebrate Community Monitoring." *Current Biology* 32 (3): 701-707.e5. <https://doi.org/10.1016/j.cub.2021.12.014>.

- Macreadie, Peter I, Ashley M Fowler, and David J Booth. 2011. "Rigs-to-Reefs: Will the Deep Sea Benefit from Artificial Habitat?" *Frontiers in Ecology and the Environment* 9 (8): 455–61. <https://doi.org/10.1890/100112>.
- Malayil, Leena, Padmini Ramachandran, Suhana Chattopadhyay, Robin Cagle, Lauren Hittle, Andrea Ottesen, Emmanuel F. Mongodin, and Amy R. Sapkota. 2020. "Metabolically-Active Bacteria in Reclaimed Water and Ponds Revealed Using Bromodeoxyuridine DNA Labeling Coupled with 16S rRNA and Shotgun Sequencing." *Water Research* 184 (October): 116185. <https://doi.org/10.1016/j.watres.2020.116185>.
- Marques, João Tiago, Maria J. Ramos Pereira, Tiago A. Marques, Carlos David Santos, Joana Santana, Pedro Beja, and Jorge M. Palmeirim. 2013. "Optimizing Sampling Design to Deal with Mist-Net Avoidance in Amazonian Birds and Bats." Edited by R. Mark Brigham. *PLoS ONE* 8 (9): e74505. <https://doi.org/10.1371/journal.pone.0074505>.
- Mason, Olivia U, Terry C Hazen, Sharon Borglin, Patrick S G Chain, Eric A Dubinsky, Julian L Fortney, James Han, et al. 2012. "Metagenome, Metatranscriptome and Single-Cell Sequencing Reveal Microbial Response to Deepwater Horizon Oil Spill." *The ISME Journal* 6 (9): 1715–27. <https://doi.org/10.1038/ismej.2012.59>.
- Mauffrey, Florian, Tristan Cordier, Laure Apothéloz-Perret-Gentil, Kristina Cermakova, Thomas Merzi, Matthieu Delefosse, Philippe Blanc, and Jan Pawlowski. 2020. "Benthic Monitoring of Oil and Gas Offshore Platforms in the North Sea Using Environmental DNA Metabarcoding." *Molecular Ecology* 30 (13): 3007–22. <https://doi.org/10.1111/mec.15698>.
- McClenaghan, Beverly, Nicole Fahner, David Cote, Julek Chawarski, Avery McCarthy, Hoda Rajabi, Greg Singer, and Mehrdad Hajibabaei. 2020. "Harnessing the Power of eDNA Metabarcoding for the Detection of Deep-Sea Fishes." Edited by Hideyuki Doi. *PLOS ONE* 15 (11): e0236540. <https://doi.org/10.1371/journal.pone.0236540>.
- McLaren, Michael R, Amy D Willis, and Benjamin J Callahan. 2019. "Consistent and Correctable Bias in Metagenomic Sequencing Experiments." *eLife* 8 (September): e46923. <https://doi.org/10.7554/eLife.46923>.
- McLean, Dianne L., Miles J. G. Parsons, Andrew R. Gates, Mark C. Benfield, Todd Bond, David J. Booth, Michael Bunce, et al. 2020. "Enhancing the Scientific Value of Industry Remotely Operated Vehicles (ROVs) in Our Oceans." *Frontiers in Marine Science* 7 (April): 220. <https://doi.org/10.3389/fmars.2020.00220>.
- Mena, José Luis, Hiromi Yagui, Vania Tejada, Emilio Bonifaz, Eva Bellemain, Alice Valentini, Mathias W. Tobler, Pamela Sánchez-Vendizú, and Arnaud Lyet. 2021. "Environmental DNA Metabarcoding as a Useful Tool for Evaluating Terrestrial Mammal Diversity in Tropical Forests." *Ecological Applications* 31 (5): e02335. <https://doi.org/10.1002/eap.2335>.
- Miya, Masaki, Ryo O. Gotoh, and Tetsuya Sado. 2020. "MiFish Metabarcoding: A High-Throughput Approach for Simultaneous Detection of Multiple Fish Species from Environmental DNA and Other Samples." *Fisheries Science* 86 (6): 939–70. <https://doi.org/10.1007/s12562-020-01461-x>.
- Montereale Gavazzi, Giacomo, Danae Athena Kapasakali, Francis Kerchhof, Samuel Deleu, Steven Degraer, and Vera Van Lancker. 2021. "Subtidal Natural Hard Substrate Quantitative Habitat Mapping: Interlinking Underwater Acoustics and Optical Imagery with Machine Learning." *Remote Sensing* 13 (22): 4608. <https://doi.org/10.3390/rs13224608>.
- Moss, Wynne E., Lynsey R. Harper, Mark A. Davis, Caren S. Goldberg, Matthew M. Smith, and Pieter T. J. Johnson. 2022. "Navigating the Trade-offs between Environmental

- DNA and Conventional Field Surveys for Improved Amphibian Monitoring.” *Ecosphere* 13 (2). <https://doi.org/10.1002/ecs2.3941>.
- Mynott, Sebastian. 2019. “Pilot Study to Validate an Environmental DNA Sampler for Monitoring Inshore Fish Communities.” NECR287. Natural England Commissioned Report.
- Nacke, Heiko, Kezia Goldmann, Ingo Schöning, Birgit Pfeiffer, Kristin Kaiser, Genis A Castillo-Villamizar, Marion Schrupf, François Buscot, Rolf Daniel, and Tesfaye Wubet. 2016. “Fine Spatial Scale Variation of Soil Microbial Communities under European Beech and Norway Spruce.” *Frontiers in Microbiology* 7: 14.
- Nakagawa, Hikaru, Satoshi Yamamoto, Yukuto Sato, Tetsuya Sado, Toshifumi Minamoto, and Masaki Miya. 2018. “Comparing Local- and Regional-scale Estimations of the Diversity of Stream Fish Using eDNA Metabarcoding and Conventional Observation Methods.” *Freshwater Biology* 63 (6): 569–80. <https://doi.org/10.1111/fwb.13094>.
- Newton, Joshua P., Philip W. Bateman, Matthew J. Heydenrych, Joshua H. Kestel, Kingsley W. Dixon, Kit S. Prendergast, Nicole E. White, and Paul Nevill. 2023. “Monitoring the Birds and the Bees: Environmental DNA Metabarcoding of Flowers Detects Plant–Animal Interactions.” *Environmental DNA*, March, edn3.399. <https://doi.org/10.1002/edn3.399>.
- Nguyen, Ngoc-Loi, Dhanushka Devendra, Natalia Szymańska, Mattia Greco, Inès Barrenechea Angeles, Agnes K. M. Weiner, Jessica Louise Ray, et al. 2023. “Sedimentary Ancient DNA: A New Paleogenomic Tool for Reconstructing the History of Marine Ecosystems.” *Frontiers in Marine Science* 10 (June): 1185435. <https://doi.org/10.3389/fmars.2023.1185435>.
- Nguyen, Tan T., Sabine K.J. Cochrane, and Bjarne Landfald. 2018. “Perturbation of Seafloor Bacterial Community Structure by Drilling Waste Discharge.” *Marine Pollution Bulletin* 129 (2): 615–22. <https://doi.org/10.1016/j.marpolbul.2017.10.039>.
- Nistal-Garcia, Alejandro, Pedro Garcia-Garcia, Jorge Garcia-Giron, Maria Borrego-Ramos, Saul Blanco, and Eloy Bécares. 2020. “DNA Metabarcoding and Morphological Methods Show Complementary Patterns in the Metacommunity Organization of Lentic Epiphytic Diatoms.” Preprint. Preprints. <https://doi.org/10.22541/au.160037721.10490381>.
- Nordstrom, Bethany, Nicola Mitchell, Margaret Byrne, and Simon Jarman. 2022. “A Review of Applications of Environmental DNA for Reptile Conservation and Management.” *Ecology and Evolution* 12 (6). <https://doi.org/10.1002/ece3.8995>.
- Norwegian Environmental Agency. 2020. “Guidelines for Environmental Monitoring of Petroleum Activities on the Norwegian Continental Shelf.”
- Obst, Matthias, Katrina Exter, A. Louise Allcock, Christos Arvanitidis, Alizz Axberg, Maria Bustamante, Ibon Cancio, et al. 2020. “A Marine Biodiversity Observation Network for Genetic Monitoring of Hard-Bottom Communities (ARMS-MBON).” *Frontiers in Marine Science* 7 (November): 572680. <https://doi.org/10.3389/fmars.2020.572680>.
- Oladi, Mahshid, Kleopatra Leontidou, Thorsten Stoeck, and Mohammad Reza Shokri. 2022. “Environmental DNA-Based Profiling of Benthic Bacterial and Eukaryote Communities along a Crude Oil Spill Gradient in a Coral Reef in the Persian Gulf.” *Marine Pollution Bulletin* 184 (November): 114143. <https://doi.org/10.1016/j.marpolbul.2022.114143>.
- Oliverio, Angela M., Huijie Gan, Kyle Wickings, and Noah Fierer. 2018. “A DNA Metabarcoding Approach to Characterize Soil Arthropod Communities.” *Soil Biology and Biochemistry* 125 (June): 37–43. <https://doi.org/10.1016/j.soilbio.2018.06.026>.
- Palmer, Kaitlin J., Kate L. Brookes, Ian M. Davies, Ewan Edwards, and Luke Rendell. 2019. “Habitat Use of a Coastal Delphinid Population Investigated Using Passive Acoustic

- Monitoring." *Aquatic Conservation: Marine and Freshwater Ecosystems* 29 (S1): 254–70. <https://doi.org/10.1002/aqc.3166>.
- Pampanin, Daniela M., Steven J. Brooks, Bjørn Einar Grøsvik, Jérémie Le Goff, Sonnich Meier, and Magne O. Sydnes. 2017. "DNA Adducts in Marine Fish as Biological Marker of Genotoxicity in Environmental Monitoring: The Way Forward." *Marine Environmental Research* 125 (April): 49–62. <https://doi.org/10.1016/j.marenvres.2017.01.002>.
- Pawlowski, Jan, Laure Apothéoz-Perret-Gentil, Elvira Mächler, and Florian Altermatt. 2020. "Environmental DNA Applications for Biomonitoring and Bioassessment in Aquatic Ecosystems." <https://doi.org/10.5167/UZH-187800>.
- Pawlowski, Jan, K. Bruce, K. Panksep, F.I. Aguirre, S. Amalfitano, L. Apothéoz-Perret-Gentil, T. Baussant, et al. 2022. "Environmental DNA Metabarcoding for Benthic Monitoring: A Review of Sediment Sampling and DNA Extraction Methods." *Science of The Total Environment* 818 (April): 151783. <https://doi.org/10.1016/j.scitotenv.2021.151783>.
- Pawlowski, Jan, Philippe Esling, Franck Lejzerowicz, Tomas Cedhagen, and Thomas A. Wilding. 2014. "Environmental Monitoring through Protist Next-Generation Sequencing Metabarcoding: Assessing the Impact of Fish Farming on Benthic Foraminifera Communities." *Molecular Ecology Resources* 14 (6): 1129–40. <https://doi.org/10.1111/1755-0998.12261>.
- Pawlowski, Jan, Mary Kelly-Quinn, Florian Altermatt, Laure Apothéoz-Perret-Gentil, Pedro Beja, Angela Boggero, Angel Borja, et al. 2018. "The Future of Biotic Indices in the Ecogenomic Era: Integrating (e)DNA Metabarcoding in Biological Assessment of Aquatic Ecosystems." *Science of The Total Environment* 637–638 (October): 1295–1310. <https://doi.org/10.1016/j.scitotenv.2018.05.002>.
- Pearman, John K., Guillem Chust, Eva Aylagas, Ernesto Villarino, James R. Watson, Anne Chenuil, Angel Borja, et al. 2020. "Pan-regional Marine Benthic Cryptobiome Biodiversity Patterns Revealed by Metabarcoding Autonomous Reef Monitoring Structures." *Molecular Ecology* 29 (24): 4882–97. <https://doi.org/10.1111/mec.15692>.
- Pereira-da-Conceicao, Lyndall, Vasco Elbrecht, Andie Hall, Andrew Briscoe, Helen Barber-James, and Benjamin Price. 2020. "Metabarcoding Unsorted Kick-samples Facilitates Macroinvertebrate-based Biomonitoring with Increased Taxonomic Resolution, While Outperforming Environmental DNA." *Environmental DNA* 3 (2): 353–71. <https://doi.org/10.1002/edn3.116>.
- Perelman, Zander E., Mizuki K. Takahashi, Emma L. Hundermark, and Carol Parnan. 2021. "An eDNA-Based Assessment of the Impact of a 207,000-Liter Gasoline Spill on Local Populations of Eastern Hellbenders (*Cryptobranchus Alleganiensis*), an Imperiled Giant Salamander." *Northeastern Naturalist* 28 (4). <https://doi.org/10.1656/045.028.0406>.
- Pérez-Burillo, Javier, Rosa Trobajo, Valentin Vasselon, Frédéric Rimet, Agnès Bouchez, and David G. Mann. 2020. "Evaluation and Sensitivity Analysis of Diatom DNA Metabarcoding for WFD Bioassessment of Mediterranean Rivers." *Science of The Total Environment* 727 (July): 138445. <https://doi.org/10.1016/j.scitotenv.2020.138445>.
- Piaggio, Antoinette J., Richard M. Engeman, Matthew W. Hopken, John S. Humphrey, Kandy L. Keacher, William E. Bruce, and Michael L. Avery. 2014. "Detecting an Elusive Invasive Species: A Diagnostic PCR to Detect Burmese Python in Florida Waters and an Assessment of Persistence of Environmental DNA." *Molecular Ecology Resources* 14 (2): 374–80. <https://doi.org/10.1111/1755-0998.12180>.
- Pierella Karlusich, Juan José, Fabien Lombard, Jean-Olivier Irisson, Chris Bowler, and Rachel A. Foster. 2022. "Coupling Imaging and Omics in Plankton Surveys: State-of-

- the-Art, Challenges, and Future Directions." *Frontiers in Marine Science* 9 (June): 878803. <https://doi.org/10.3389/fmars.2022.878803>.
- Pierella Karlusich, Juan José, Eric Pelletier, Lucie Zinger, Fabien Lombard, Adriana Zingone, Sébastien Colin, Josep M. Gasol, et al. 2022. "A Robust Approach to Estimate Relative Phytoplankton Cell Abundances from Metagenomes." *Molecular Ecology Resources*, February, 1755-0998.13592. <https://doi.org/10.1111/1755-0998.13592>.
- Piper, Alexander M, Jana Batovska, Noel O I Cogan, John Weiss, John Paul Cunningham, Brendan C Rodoni, and Mark J Blacket. 2019. "Prospects and Challenges of Implementing DNA Metabarcoding for High-Throughput Insect Surveillance." *GigaScience* 8 (8): giz092. <https://doi.org/10.1093/gigascience/giz092>.
- Pitois, Sophie G., Carolyn A. Graves, Hayden Close, Christopher Lynam, James Scott, Julian Tilbury, Jeroen van der Kooij, and Phil Culverhouse. 2021. "A First Approach to Build and Test the Copepod Mean Size and Total Abundance (CMSTA) Ecological Indicator Using in-Situ Size Measurements from the Plankton Imager (PI)." *Ecological Indicators* 123 (April): 107307. <https://doi.org/10.1016/j.ecolind.2020.107307>.
- Polanco Fernández, Andrea, Virginie Marques, Fabian Fopp, Jean Baptiste Juhel, Giomar Helena Borrero-Pérez, Marie Charlotte Cheutin, Tony Dejean, et al. 2021. "Comparing Environmental DNA Metabarcoding and Underwater Visual Census to Monitor Tropical Reef Fishes." *Environmental DNA* 3 (1): 142–56. <https://doi.org/10.1002/edn3.140>.
- Polanco Fernández, Andrea, Maria Mutis Martinezguerra, Virginie Marques, Francisco Villa-Navarro, Giomar Helena Borrero Pérez, Marie-Charlotte Cheutin, Tony Dejean, et al. 2021. "Detecting Aquatic and Terrestrial Biodiversity in a Tropical Estuary Using Environmental DNA." *Biotropica* 53 (6): 1606–19. <https://doi.org/10.1111/btp.13009>.
- Pollina, Thibaut, Adam G. Larson, Fabien Lombard, Hongquan Li, David Le Guen, Sébastien Colin, Colomban De Vargas, and Manu Prakash. 2022. "PlanktoScope: Affordable Modular Quantitative Imaging Platform for Citizen Oceanography." *Frontiers in Marine Science* 9 (July): 949428. <https://doi.org/10.3389/fmars.2022.949428>.
- Pont, Didier, Paul Meulenbroek, Vincenz Bammer, Tony Dejean, Tibor Erős, Pauline Jean, Mirjana Lenhardt, et al. 2022. "Quantitative Monitoring of Diverse Fish Communities on a Large Scale Combining eDNA Metabarcoding and qPCR." *Molecular Ecology Resources*, October, 1755-0998.13715. <https://doi.org/10.1111/1755-0998.13715>.
- Pont, Didier, Mathieu Rocle, Alice Valentini, Raphaël Civade, Pauline Jean, Anthony Maire, Nicolas Roset, Michael Schabuss, Horst Zornig, and Tony Dejean. 2018. "Environmental DNA Reveals Quantitative Patterns of Fish Biodiversity in Large Rivers despite Its Downstream Transportation." *Scientific Reports* 8 (1): 10361. <https://doi.org/10.1038/s41598-018-28424-8>.
- Port, Jesse A., James L. O'Donnell, Ofelia C. Romero-Maraccini, Paul R. Leary, Steven Y. Litvin, Kerry J. Nickols, Kevan M. Yamahara, and Ryan P. Kelly. 2016. "Assessing Vertebrate Biodiversity in a Kelp Forest Ecosystem Using Environmental DNA." *Molecular Ecology* 25 (2): 527–41. <https://doi.org/10.1111/mec.13481>.
- Pukk, Lilian, Jeannette Kanefsky, Amanda L. Heathman, Ellen M. Weise, Lucas R. Nathan, Seth J. Herbst, Nicholas M. Sard, Kim T. Scribner, and John D. Robinson. 2021. "eDNA Metabarcoding in Lakes to Quantify Influences of Landscape Features and Human Activity on Aquatic Invasive Species Prevalence and Fish Community Diversity." Edited by Catherine Richter. *Diversity and Distributions* 27 (10): 2016–31. <https://doi.org/10.1111/ddi.13370>.
- Rees, Helen C., Keith Bishop, David J. Middleditch, James R. M. Patmore, Ben C. Maddison, and Kevin C. Gough. 2014. "The Application of eDNA for Monitoring of the

- Great Crested Newt in the UK." *Ecology and Evolution* 4 (21): 4023–32. <https://doi.org/10.1002/ece3.1272>.
- Riaz, Maria, Mathias Kuemmerlen, Claudia Wittwer, Berardino Cocchiararo, Imran Khaliq, Markus Pfenninger, and Carsten Nowak. 2020. "Combining Environmental DNA and Species Distribution Modeling to Evaluate Reintroduction Success of a Freshwater Fish." *Ecological Applications* 30 (2). <https://doi.org/10.1002/eap.2034>.
- Rice, Christopher J., Eric R. Larson, and Christopher A. Taylor. 2018. "Environmental DNA Detects a Rare Large River Crayfish but with Little Relation to Local Abundance." *Freshwater Biology* 63 (5): 443–55. <https://doi.org/10.1111/fwb.13081>.
- Rimet, Frédéric, and A. Bouchez. 2012. "Life-Forms, Cell-Sizes and Ecological Guilds of Diatoms in European Rivers." *Knowledge and Management of Aquatic Ecosystems*, no. 406: 01. <https://doi.org/10.1051/kmae/2012018>.
- Rimet, Frédéric, Evgeny Gusev, Maria Kahlert, Martyn G. Kelly, Maxim Kulikovskiy, Yevhen Maltsev, David G. Mann, et al. 2019. "Diat.Barcode, an Open-Access Curated Barcode Library for Diatoms." *Scientific Reports* 9 (1): 15116. <https://doi.org/10.1038/s41598-019-51500-6>.
- Rivera, Sinziana F., Valentin Vasselon, Agnès Bouchez, and Frédéric Rimet. 2023. "eDNA Metabarcoding from Aquatic Biofilms Allows Studying Spatial and Temporal Fluctuations of Fish Communities from Lake Geneva." *Environmental DNA* 5 (3): 570–81. <https://doi.org/10.1002/edn3.413>.
- Rivera, Sinziana F., Valentin Vasselon, Nathalie Mary, Olivier Monnier, Frédéric Rimet, and Agnès Bouchez. 2021. "Exploring the Capacity of Aquatic Biofilms to Act as Environmental DNA Samplers: Test on Macroinvertebrate Communities in Rivers." *Science of The Total Environment* 763 (April): 144208. <https://doi.org/10.1016/j.scitotenv.2020.144208>.
- Robinson, Jake M., Peter A. Harrison, Suzanne Mavoia, and Martin F. Breed. 2022. "Existing and Emerging Uses of Drones in Restoration Ecology." *Methods in Ecology and Evolution* 13 (9): 1899–1911. <https://doi.org/10.1111/2041-210X.13912>.
- Roger, Fabian, Hamid Ghanavi, Natalie Danielsson, Niklas Wahlberg, Jakob Löndahl, Lars B. Pettersson, Georg K.S. Andersson, Niklas Boke Olén, and Yann Clough. 2022. "Airborne Environmental DNA Metabarcoding for the Monitoring of Terrestrial Insects - a Proof of Concept," *Environmental DNA*, 4 (4): 790–807. <https://doi.org/10.1101/2021.07.26.453860>.
- Rojahn, Jack, Dianne Gleeson, and Elise M. Furlan. 2018. "Monitoring Post-Release Survival of the Northern Corroboree Frog, *Pseudophryne pengillyi*, Using Environmental DNA." *Wildlife Research* 45 (7): 620. <https://doi.org/10.1071/WR17179>.
- Rosa, Luiz Henrique, Tamar Holanda da Silva, Mayara Baptistucci Ogaki, Otávio Henrique Bezerra Pinto, Michael Stech, Peter Convey, Micheline Carvalho-Silva, Carlos Augusto Rosa, and Paulo E. A. S. Câmara. 2020. "DNA Metabarcoding Uncovers Fungal Diversity in Soils of Protected and Non-Protected Areas on Deception Island, Antarctica." *Scientific Reports* 10 (1): 21986. <https://doi.org/10.1038/s41598-020-78934-7>.
- Rose, Jonathan P., Cara Wademan, Suzanne Weir, John S. Wood, and Brian D. Todd. 2019. "Traditional Trapping Methods Outperform eDNA Sampling for Introduced Semi-Aquatic Snakes." Edited by Benedikt R. Schmidt. *PLOS ONE* 14 (7): e0219244. <https://doi.org/10.1371/journal.pone.0219244>.
- Rosenberg, Rutger, Mats Blomqvist, Hans C Nilsson, Hans Cederwall, and Anna Dimming. 2004. "Marine Quality Assessment by Use of Benthic Species-Abundance Distributions: A Proposed New Protocol within the European Union Water Framework

- Directive." *Marine Pollution Bulletin* 49 (9–10): 728–39.
<https://doi.org/10.1016/j.marpolbul.2004.05.013>.
- Roux, Louise-Marie D., Danièle Giblot-Ducray, Nathan J. Bott, Kathryn H. Wiltshire, Marty R. Deveney, Kristen M. Westfall, and Cathryn L. Abbott. 2020. "Analytical Validation and Field Testing of a Specific qPCR Assay for Environmental DNA Detection of Invasive European Green Crab (*Carcinus Maenas*)." *Environmental DNA* 2 (3): 309–20.
<https://doi.org/10.1002/edn3.65>.
- Ryan, Ethan, Phil Bateman, Mieke van der Heyde, Kristen Fernandes, and Paul Nevill. 2020. "Comparison of eDNA Metabarcoding to Camera Trapping for Terrestrial Vertebrate Monitoring Highlights the Importance of Substrate Type, Frequency of Sampling and Animal Size." Preprint. Preprints.
<https://doi.org/10.22541/au.160442291.13614413/v1>.
- Ryan, Ethan, Philip Bateman, Kristen Fernandes, Mieke Heyde, and Paul Nevill. 2022. "EDNA Metabarcoding of Log Hollow Sediments and Soils Highlights the Importance of Substrate Type, Frequency of Sampling and Animal Size, for Vertebrate Species Detection." *Environmental DNA*, May, edn3.306. <https://doi.org/10.1002/edn3.306>.
- Rzeznik-Orignac, Jadwiga, Dimitri Kalenitchenko, Jérôme Mariette, Jean-Yves Bodiou, Nadine Le Bris, and Evelyne Derelle. 2017. "Comparison of Meiofaunal Diversity by Combined Morphological and Molecular Approaches in a Shallow Mediterranean Sediment." *Marine Biology* 164 (3): 40. <https://doi.org/10.1007/s00227-017-3074-4>.
- Sakata, Masayuki K., Satoshi Yamamoto, Ryo O. Gotoh, Masaki Miya, Hiroki Yamanaka, and Toshifumi Minamoto. 2020. "Sedimentary eDNA Provides Different Information on Timescale and Fish Species Composition Compared with Aqueous eDNA." *Environmental DNA* 2 (4): 505–18. <https://doi.org/10.1002/edn3.75>.
- Sales, Naiara Guimarães, Maisie B. McKenzie, Joseph Drake, Lynsey R. Harper, Samuel S. Browett, Ilaria Coscia, Owen S. Wangensteen, et al. 2020. "Fishing for Mammals: Landscape-level Monitoring of Terrestrial and Semi-aquatic Communities Using eDNA from Riverine Systems." Edited by Brittany Mosher. *Journal of Applied Ecology* 57 (4): 707–16. <https://doi.org/10.1111/1365-2664.13592>.
- Schabacker, Jenna C., Stephen J. Amish, Bonnie K. Ellis, Beth Gardner, Dulaney L. Miller, Elizabeth A. Rutledge, Adam J. Sepulveda, and Gordon Luikart. 2020. "Increased eDNA Detection Sensitivity Using a Novel High-volume Water Sampling Method." *Environmental DNA* 2 (2): 244–51. <https://doi.org/10.1002/edn3.63>.
- Schenk, Janina, Sebastian Höss, Marvin Brinke, Nils Kleinbölting, Henrike Brüchner-Hüttemann, and Walter Traunspurger. 2020. "Nematodes as Bioindicators of Polluted Sediments Using Metabarcoding and Microscopic Taxonomy." *Environment International* 143 (October): 105922. <https://doi.org/10.1016/j.envint.2020.105922>.
- Schneider, Stefan, Saul Greenberg, Graham W. Taylor, and Stefan C. Kremer. 2020. "Three Critical Factors Affecting Automated Image Species Recognition Performance for Camera Traps." *Ecology and Evolution* 10 (7): 3503–17.
<https://doi.org/10.1002/ece3.6147>.
- Scholin, Christopher, James Birch, Scott Jensen, Roman Marin III, Eugene Massion, Douglas Pargett, Christina Preston, Brent Roman, William Ussler III, and William Ussler III. 2017. "The Quest to Develop Ecogenomic Sensors: A 25-Year History of the Environmental Sample Processor (ESP) as a Case Study." *Oceanography* 30 (4): 100–113. <https://doi.org/10.5670/oceanog.2017.427>.
- Searcy, Ryan T., Alexandria B. Boehm, Chloe Weinstock, Christina M. Preston, Scott Jensen, Brent Roman, James M. Birch, et al. 2022. "High-Frequency and Long-Term Observations of eDNA from Imperiled Salmonids in a Coastal Stream: Temporal Dynamics, Relationships with Environmental Factors, and Comparisons with

- Conventional Observations." *Environmental DNA* 4 (4): 776–89. <https://doi.org/10.1002/edn3.293>.
- Seymour, Mathew, Francois K Edwards, Bernard J Cosby, Martyn G Kelly, Mark de Bruyn, Gary R Carvalho, and Simon Creer. 2020. "Executing Multi-Taxa eDNA Ecological Assessment via Traditional Metrics and Interactive Networks." *Science of The Total Environment* 729 (August): 138801. <https://doi.org/10.1016/j.scitotenv.2020.138801>.
- Shelton, Andrew Olaf, Zachary J Gold, Alexander J Jensen, Erin D'Agnese, Elizabeth Andruszkiewicz, and P Kelly. 2022. "Toward Quantitative Metabarcoding," 31.
- Shu, Lu, Shijing Chen, Ping Li, and Zuogang Peng. 2022. "Environmental DNA Metabarcoding Reflects Fish DNA Dynamics in Lentic Ecosystems: A Case Study of Freshwater Ponds." *Fishes* 7 (5): 257. <https://doi.org/10.3390/fishes7050257>.
- Siano, Raffaele, Malwenn Lassudrie, Pierre Cuzin, Nicolas Briant, Véronique Loizeau, Sabine Schmidt, Axel Ehrhold, et al. 2021. "Sediment Archives Reveal Irreversible Shifts in Plankton Communities after World War II and Agricultural Pollution." *Current Biology* 31 (12): 2682-2689.e7. <https://doi.org/10.1016/j.cub.2021.03.079>.
- Silverman, Justin D., Rachael J. Bloom, Sharon Jiang, Heather K. Durand, Eric Dallow, Sayan Mukherjee, and Lawrence A. David. 2021. "Measuring and Mitigating PCR Bias in Microbiota Datasets." Edited by Alice Carolyn McHardy. *PLOS Computational Biology* 17 (7): e1009113. <https://doi.org/10.1371/journal.pcbi.1009113>.
- Slade, Norman A., and Susan M. Blair. 2000. "AN EMPIRICAL TEST OF USING COUNTS OF INDIVIDUALS CAPTURED AS INDICES OF POPULATION SIZE." *Journal of Mammalogy* 81 (4): 1035–45. [https://doi.org/10.1644/1545-1542\(2000\)081<1035:AETOUCC>2.0.CO;2](https://doi.org/10.1644/1545-1542(2000)081<1035:AETOUCC>2.0.CO;2).
- Snyder, Darrel E. 2003. "Invited Overview: Conclusions from a Review of Electrofishing and Its Harmful Effects on Fish." *Reviews in Fish Biology and Fisheries* 13 (4): 445–53. <https://doi.org/10.1007/s11160-004-1095-9>.
- Snyder, Matthew R, and Carol A. Stepien. 2020. "Increasing Confidence for Discerning Species and Population Compositions from Metabarcoding Assays of Environmental Samples: Case Studies of Fishes in the Laurentian Great Lakes and Wabash River." *Metabarcoding and Metagenomics* 4: 18.
- Song, Chi-une, Hyeongwoo Choi, Min-Seung Jeon, Eun-Jeong Kim, Hyeon Gyeong Jeong, Sung Kim, Choong-gon Kim, et al. 2021. "Zooplankton Diversity Monitoring Strategy for the Urban Coastal Region Using Metabarcoding Analysis." *Scientific Reports* 11 (1): 24339. <https://doi.org/10.1038/s41598-021-03656-3>.
- Spence, Brian C., David E. Rundio, Nicholas J. Demetras, and Maryna Sedoryk. 2021. "Efficacy of Environmental DNA Sampling to Detect the Occurrence of Endangered Coho Salmon (*Oncorhynchus Kisutch*) in Mediterranean-climate Streams of California's Central Coast." *Environmental DNA* 3 (4): 727–44. <https://doi.org/10.1002/edn3.175>.
- Spotswood, Erica N., Kari Roesch Goodman, Jay Carlisle, Renée L. Cormier, Diana L. Humple, Josée Rousseau, Susan L. Guers, and Gina G. Barton. 2012. "How Safe Is Mist Netting? Evaluating the Risk of Injury and Mortality to Birds: *How Safe Is Mist Netting?*" *Methods in Ecology and Evolution* 3 (1): 29–38. <https://doi.org/10.1111/j.2041-210X.2011.00123.x>.
- Stefani, Franck, Nathalie Isabel, Marie-Josée Morency, Manuel Lamothe, Simon Nadeau, Denis Lachance, Edith H. Y. Li, et al. 2018. "The Impact of Reconstructed Soils Following Oil Sands Exploitation on Aspen and Its Associated Belowground Microbiome." *Scientific Reports* 8 (1): 2761. <https://doi.org/10.1038/s41598-018-20783-6>.

- Stefani, Franck O. P., Terrence H. Bell, Charlotte Marchand, Ivan E. de la Providencia, Abdel El Yassimi, Marc St-Arnaud, and Mohamed Hijri. 2015. "Culture-Dependent and -Independent Methods Capture Different Microbial Community Fractions in Hydrocarbon-Contaminated Soils." Edited by Shuijin Hu. *PLOS ONE* 10 (6): e0128272. <https://doi.org/10.1371/journal.pone.0128272>.
- Steyaert, Margaux, Victoria Priestley, Owen Osborne, Alba Herraiz, Richard Arnold, and Vincent Savolainen. 2020. "Advances in Metabarcoding Techniques Bring Us Closer to Reliable Monitoring of the Marine Benthos." Edited by Vitor Paiva. *Journal of Applied Ecology* 57 (11): 2234–45. <https://doi.org/10.1111/1365-2664.13729>.
- Stoeckle, Mark Y., Jesse H. Ausubel, and Michael Coogan. 2022. "12S Gene Metabarcoding with DNA Standard Quantifies Marine Bony Fish Environmental DNA, Identifies Threshold for Reproducible Amplification, and Overcomes Distortion Due to Non-Fish Vertebrate DNA." *bioRxiv*. <https://doi.org/10.1101/2022.07.29.502053>.
- Sugai, Larissa Sayuri Moreira, Thiago Sanna Freire Silva, José Wagner Ribeiro, and Diego Llusia. 2019. "Terrestrial Passive Acoustic Monitoring: Review and Perspectives." *BioScience* 69 (1): 15–25. <https://doi.org/10.1093/biosci/biy147>.
- Suter, Leonie, Andrea Maree Polanowski, Laurence John Clarke, John Andrew Kitchener, and Bruce Emerson Deagle. 2021. "Capturing Open Ocean Biodiversity: Comparing Environmental DNA Metabarcoding to the Continuous Plankton Recorder." *Molecular Ecology* 30 (13): 3140–57. <https://doi.org/10.1111/mec.15587>.
- Svenningsen, Anne Katrine Nørgaard, Cino Pertoldi, and Dan Bruhn. 2022. "eDNA Metabarcoding Benchmarked towards Conventional Survey Methods in Amphibian Monitoring." *Animals* 12 (6): 763. <https://doi.org/10.3390/ani12060763>.
- Taylor, J. C, W. R Harding, C. G. M Archibald, South Africa, and Water Research Commission. 2007. *A Methods Manual for the Collection, Preparation and Analysis of Diatom Samples, Version 1.0*. Gezina, South Africa: Water Research Commission.
- Tegally, Houriiyah, James Emmanuel San, Jennifer Giandhari, and Tulio de Oliveira. 2020. "Unlocking the Efficiency of Genomics Laboratories with Robotic Liquid-Handling." *BMC Genomics* 21 (1): 729. <https://doi.org/10.1186/s12864-020-07137-1>.
- Thalinger, Bettina, Kristy Deiner, Lynsey R. Harper, Helen C. Rees, Rosetta C. Blackman, Daniela Sint, Michael Traugott, Caren S. Goldberg, and Kat Bruce. 2021. "A Validation Scale to Determine the Readiness of Environmental DNA Assays for Routine Species Monitoring." *Environmental DNA* 3 (4): 823–36. <https://doi.org/10.1002/edn3.189>.
- The eDNA Society. 2019. "Environmental DNA Sampling and Experiment Manual Version 2.1." Otsu, Japan.
- Thomas, Austen C., Samantha Tank, Phong L. Nguyen, Jake Ponce, Mieke Sinnesael, and Caren S. Goldberg. 2020. "A System for Rapid eDNA Detection of Aquatic Invasive Species." *Environmental DNA* 2 (3): 261–70. <https://doi.org/10.1002/edn3.25>.
- Thomas, Morgan L., Lynn Baker, James R. Beattie, and Andrew M. Baker. 2020. "Determining the Efficacy of Camera Traps, Live Capture Traps, and Detection Dogs for Locating Cryptic Small Mammal Species." *Ecology and Evolution* 10 (2): 1054–68. <https://doi.org/10.1002/ece3.5972>.
- Thompson, Graham G., and Scott A. Thompson. 2007. "Usefulness of Funnel Traps in Catching Small Reptiles and Mammals, with Comments on the Effectiveness of the Alternatives." *Wildlife Research* 34 (6): 491. <https://doi.org/10.1071/WR06081>.
- Thomsen, Philip Francis, Peter Rask Møller, Eva Egelyng Sigsgaard, Steen Wilhelm Knudsen, Ole Ankjær Jørgensen, and Eske Willerslev. 2016. "Environmental DNA from Seawater Samples Correlate with Trawl Catches of Subarctic, Deepwater

- Fishes." Edited by Andrew R. Mahon. *PLOS ONE* 11 (11): e0165252. <https://doi.org/10.1371/journal.pone.0165252>.
- Tiburcio, Samyra Raquel Gonçalves, Andrew Macrae, Raquel Silva Peixoto, Caio Tavora Coelho da Costa Rachid, Felipe Raposo Passos Mansoldo, Daniela Sales Alviano, Celuta Sales Alviano, et al. 2021. "Sulphate-Reducing Bacterial Community Structure from Produced Water of the Periquito and Galo de Campina Onshore Oilfields in Brazil." *Scientific Reports* 11 (1): 20311. <https://doi.org/10.1038/s41598-021-99196-x>.
- Todd, Jacqui H., Robert M. Simpson, Joanne Poulton, Emma I. Barraclough, Kurt Villsen, Amber Brooks, Kate Richards, and Dan Jones. 2020. "Detecting Invertebrate Ecosystem Service Providers in Orchards: Traditional Methods versus Barcoding of Environmental DNA in Soil." *Agricultural and Forest Entomology* 22 (3): 212–23. <https://doi.org/10.1111/afe.12374>.
- Torre, Ignasi, Lúcia Freixas, Antoni Arrizabalaga, and Mario Díaz. 2016. "The Efficiency of Two Widely Used Commercial Live-Traps to Develop Monitoring Protocols for Small Mammal Biodiversity." *Ecological Indicators* 66 (July): 481–87. <https://doi.org/10.1016/j.ecolind.2016.02.017>.
- Treonis, Amy M., Samantha K. Unangst, Ryan M. Kepler, Jeffrey S. Buyer, Michel A. Cavigelli, Steven B. Mirsky, and Jude E. Maul. 2018. "Characterization of Soil Nematode Communities in Three Cropping Systems through Morphological and DNA Metabarcoding Approaches." *Scientific Reports* 8 (1): 2004. <https://doi.org/10.1038/s41598-018-20366-5>.
- Trevelin, Leonardo Carreira, Roberto Leonan Morim Novaes, Paul François Colas-Rosas, Thayse Cristhina Melo Benathar, and Carlos A. Peres. 2017. "Enhancing Sampling Design in Mist-Net Bat Surveys by Accounting for Sample Size Optimization." Edited by Danilo Russo. *PLOS ONE* 12 (3): e0174067. <https://doi.org/10.1371/journal.pone.0174067>.
- Trexler, Ryan, Caroline Solomon, Colin J. Brislawn, Justin R. Wright, Abigail Rosenberger, Erin E. McClure, Alyssa M. Grube, et al. 2014. "Assessing Impacts of Unconventional Natural Gas Extraction on Microbial Communities in Headwater Stream Ecosystems in Northwestern Pennsylvania." *Frontiers in Microbiology* 5 (November). <https://doi.org/10.3389/fmicb.2014.00522>.
- Truelove, Nathan K., Nastassia V. Patin, Markus Min, Kathleen J. Pitz, Chris M. Preston, Kevan M. Yamahara, Yanwu Zhang, et al. 2022. "Expanding the Temporal and Spatial Scales of Environmental DNA Research with Autonomous Sampling." *Environmental DNA* 4 (4): 972–84. <https://doi.org/10.1002/edn3.299>.
- Ulrich, Nikea, Veronica Kirchner, Rebecca Drucker, Justin R. Wright, Christopher J. McLimans, Terry C. Hazen, Maria F. Campa, Christopher J. Grant, and Regina Lamendella. 2018. "Response of Aquatic Bacterial Communities to Hydraulic Fracturing in Northwestern Pennsylvania: A Five-Year Study." *Scientific Reports* 8 (1): 5683. <https://doi.org/10.1038/s41598-018-23679-7>.
- Urban, Lara, Andre Holzer, J Jotautas Baronas, Michael B Hall, Philipp Braeuninger-Weimer, Michael J Scherm, Daniel J Kunz, et al. 2021. "Freshwater Monitoring by Nanopore Sequencing." *eLife* 10 (January): e61504. <https://doi.org/10.7554/eLife.61504>.
- Ushio, Masayuki, Hisato Fukuda, Toshiki Inoue, Kobayashi Makoto, Osamu Kishida, Keiichi Sato, Koichi Murata, et al. 2017. "Environmental DNA Enables Detection of Terrestrial Mammals from Forest Pond Water." *Molecular Ecology Resources* 17 (6). <https://doi.org/10.1111/1755-0998.12690>.
- Valdivia-Carrillo, Tania, Axayácatl Rocha-Olivares, Héctor Reyes-Bonilla, José Francisco Domínguez-Contreras, and Adrian Munguia-Vega. 2021. "Integrating eDNA Metabarcoding and Simultaneous Underwater Visual Surveys to Describe Complex

- Fish Communities in a Marine Biodiversity Hotspot." *Molecular Ecology Resources* 21 (5): 1558–74. <https://doi.org/10.1111/1755-0998.13375>.
- Valentin, Rafael E., Dina M. Fonseca, Sydney Gable, Kathleen E. Kyle, George C. Hamilton, Anne L. Nielsen, and Julie L. Lockwood. 2020. "Moving eDNA Surveys onto Land: Strategies for Active eDNA Aggregation to Detect Invasive Forest Insects." *Molecular Ecology Resources* 20 (3): 746–55. <https://doi.org/10.1111/1755-0998.13151>.
- Valentini, Alice, Pierre Taberlet, Claude Miaud, Raphaël Civade, Jelger Herder, Philip Francis Thomsen, Eva Bellemain, et al. 2016. "Next-Generation Monitoring of Aquatic Biodiversity Using Environmental DNA Metabarcoding." *Molecular Ecology* 25 (4): 929–42. <https://doi.org/10.1111/mec.13428>.
- Valsecchi, Elena, Jonas Bylemans, Simon J. Goodman, Roberto Lombardi, Ian Carr, Laura Castellano, Andrea Galimberti, and Paolo Galli. 2020. "Novel Universal Primers for Metabarcoding Environmental DNA Surveys of Marine Mammals and Other Marine Vertebrates." *Environmental DNA* 2 (4): 460–76. <https://doi.org/10.1002/edn3.72>.
- Vasselon, Valentin, Éva Ács, Salomé Almeida, Karl Andree, Laure Apothéoz-Perret-Gentil, Bonnie Baillet, Ana Baricevic, et al. 2021. "The Fellowship of the Ring Test: DNAqua-Net WG2 Initiative to Compare Diatom Metabarcoding Protocols Used in Routine Freshwater Biomonitoring for Standardisation." *ARPHA Conference Abstracts 4* (March): e65142. <https://doi.org/10.3897/aca.4.e65142>.
- Vasselon, Valentin, Frédéric Rimet, Kálmán Tapolczai, and Agnès Bouchez. 2017. "Assessing Ecological Status with Diatoms DNA Metabarcoding: Scaling-up on a WFD Monitoring Network (Mayotte Island, France)." *Ecological Indicators* 82 (November): 1–12. <https://doi.org/10.1016/j.ecolind.2017.06.024>.
- Verdier, Héloïse, Lara Konecny, Christophe Marquette, Helen Reveron, Laurent Grémillard, Amélie Barthès, Thibault Datry, Agnès Bouchez, and Tristan Lefébure. 2022. "Passive Sampling of Environmental DNA in Aquatic Environments Using 3D-Printed Hydroxyapatite Samplers." *Molecular Ecology Resources* 22 (6).
- Visco, Joana Amorim, Laure Apothéoz-Perret-Gentil, Arielle Cordonier, Philippe Esling, Loïc Pillet, and Jan Pawlowski. 2015. "Environmental Monitoring: Inferring the Diatom Index from Next-Generation Sequencing Data." *Environmental Science & Technology* 49 (13): 7597–7605. <https://doi.org/10.1021/es506158m>.
- Vivien, Régis, Laure Apothéoz-Perret-Gentil, Jan Pawlowski, Inge Werner, Michel Lafont, and Benoit J. D. Ferrari. 2020. "High-Throughput DNA Barcoding of Oligochaetes for Abundance-Based Indices to Assess the Biological Quality of Sediments in Streams and Lakes." *Scientific Reports* 10 (1): 2041. <https://doi.org/10.1038/s41598-020-58703-2>.
- Vivien, Régis, Franck Lejzerowicz, and Jan Pawlowski. 2016. "Next-Generation Sequencing of Aquatic Oligochaetes: Comparison of Experimental Communities." Edited by Peter Prentis. *PLOS ONE* 11 (2): e0148644. <https://doi.org/10.1371/journal.pone.0148644>.
- Wacker, Sebastian, Frode Fossøy, Bjørn Mejdell Larsen, Hege Brandsegg, Rolf Sivertsgård, and Sten Karlsson. 2019. "Downstream Transport and Seasonal Variation in Freshwater Pearl Mussel (*Margaritifera Margaritifera*) eDNA Concentration." *Environmental DNA* 1 (1): 64–73. <https://doi.org/10.1002/edn3.10>.
- Wang, Hui, Caixia Wang, Meng Lin, Xiangnan Sun, Chuanyuan Wang, and Xiaoke Hu. 2013. "Phylogenetic Diversity of Bacterial Communities Associated with Bioremediation of Crude Oil in Microcosms." *International Biodeterioration & Biodegradation* 85 (November): 400–406. <https://doi.org/10.1016/j.ibiod.2013.07.015>.
- Wangensteen, Owen S., Emma Cebrian, Creu Palacín, and Xavier Turon. 2018. "Under the Canopy: Community-Wide Effects of Invasive Algae in Marine Protected Areas Revealed by Metabarcoding." *Marine Pollution Bulletin* 127 (February): 54–66. <https://doi.org/10.1016/j.marpolbul.2017.11.033>.

- Wangensteen, Owen S., Creu Palacín, Magdalena Guardiola, and Xavier Turon. 2018. "DNA Metabarcoding of Littoral Hard-Bottom Communities: High Diversity and Database Gaps Revealed by Two Molecular Markers." *PeerJ* 6 (May): e4705. <https://doi.org/10.7717/peerj.4705>.
- Wani, Atif Khurshid, Nahid Akhtar, Nafiaah Naqash, Chirag Chopra, Reena Singh, Vineet Kumar, Sunil Kumar, Sikandar I. Mulla, and Juliana Heloisa Pinê Américo-Pinheiro. 2022. "Bioprospecting Culturable and Unculturable Microbial Consortia through Metagenomics for Bioremediation." *Cleaner Chemical Engineering* 2 (June): 100017. <https://doi.org/10.1016/j.clce.2022.100017>.
- Watts, Corinne, Andrew Dopheide, Robert Holdaway, Carina Davis, Jamie Wood, Danny Thornburrow, and Ian A Dickie. 2019. "DNA Metabarcoding as a Tool for Invertebrate Community Monitoring: A Case Study Comparison with Conventional Techniques." *Austral Entomology* 58 (3): 675–86. <https://doi.org/10.1111/aen.12384>.
- West, Katrina M., Matthew Heydenrych, Rose Lines, Tony Tucker, Sabrina Fossette, Scott Whiting, and Michael Bunce. 2021. "Development of a 16S Metabarcoding Assay for the Environmental DNA (eDNA) Detection of Aquatic Reptiles across Northern Australia." *Marine and Freshwater Research*. <https://doi.org/10.1071/MF20288>.
- Wevers, Jolien, Natalie Beenaerts, Jim Casaer, Fridolin Zimmermann, Tom Artois, and Julien Fattebert. 2021. "Modelling Species Distribution from Camera Trap By-catch Using a Scale-optimized Occupancy Approach." Edited by Marcus Rowcliffe and Rahel Sollmann. *Remote Sensing in Ecology and Conservation* 7 (3): 534–49. <https://doi.org/10.1002/rse2.207>.
- Whitaker, Justine M., Amy L. Brower, and Alexis M. Janosik. 2021. "Invasive Lionfish Detected in Estuaries in the Northern Gulf of Mexico Using Environmental DNA." *Environmental Biology of Fishes* 104 (11): 1475–85. <https://doi.org/10.1007/s10641-021-01177-6>.
- Williams, Kelly E., Kathryn P. Huyvaert, Kurt C. Vercauteren, Amy J. Davis, and Antoinette J. Piaggio. 2018. "Detection and Persistence of Environmental DNA from an Invasive, Terrestrial Mammal." *Ecology and Evolution* 8 (1): 688–95. <https://doi.org/10.1002/ece3.3698>.
- Williams, Molly-Ann, Joyce O'Grady, Bernard Ball, Jens Carlsson, Elvira Eyto, Philip McGinnity, Eleanor Jennings, Fiona Regan, and Anne Parle-McDermott. 2019. "The Application of CRISPR-Cas for Single Species Identification from Environmental DNA." *Molecular Ecology Resources* 19 (5): 1106–14. <https://doi.org/10.1111/1755-0998.13045>.
- Wilson, Deborah J, Robin L Mulvey, and Ryan D Clark. 2007. "Sampling Skinks and Geckos in Artificial Cover Objects in a Dry Mixed." *NEW ZEALAND JOURNAL OF ECOLOGY* 31 (2).
- Wort, E, H Flinham, E Good, J Bakker, and H & Egeter B Craig. 2022. "Review of DNA-Based Marine Benthic Monitoring Protocols (JNCC Report No. 705)," 41.
- Xie, Yuwei, Xiaowei Zhang, Jianghua Yang, Seonjin Kim, Seongjin Hong, John P. Giesy, Un Hyuk Yim, Won Joon Shim, Hongxia Yu, and Jong Seong Khim. 2018. "eDNA-Based Bioassessment of Coastal Sediments Impacted by an Oil Spill." *Environmental Pollution* 238 (July): 739–48. <https://doi.org/10.1016/j.envpol.2018.02.081>.
- Yamahara, Kevan M., Christina M. Preston, James Birch, Kristine Walz, Roman Marin, Scott Jensen, Douglas Pargett, et al. 2019a. "In Situ Autonomous Acquisition and Preservation of Marine Environmental DNA Using an Autonomous Underwater Vehicle." *Frontiers in Marine Science* 6 (JUL): 1–14. <https://doi.org/10.3389/fmars.2019.00373>.

- . 2019b. “In Situ Autonomous Acquisition and Preservation of Marine Environmental DNA Using an Autonomous Underwater Vehicle.” *Frontiers in Marine Science* 6 (July): 373. <https://doi.org/10.3389/fmars.2019.00373>.
- Yamamoto, Satoshi, Reiji Masuda, Yukuto Sato, Tetsuya Sado, Hitoshi Araki, Michio Kondoh, Toshifumi Minamoto, and Masaki Miya. 2017. “Environmental DNA Metabarcoding Reveals Local Fish Communities in a Species-Rich Coastal Sea.” *Scientific Reports* 7 (1): 40368. <https://doi.org/10.1038/srep40368>.
- Yang, Jianghua, Lijuan Zhang, Yawen Mu, and Xiaowei Zhang. 2023. “Small Changes Make Big Progress: A More Efficient eDNA Monitoring Method for Freshwater Fish.” *Environmental DNA*, January, edn3.387. <https://doi.org/10.1002/edn3.387>.
- Yergeau, Etienne, Christine Maynard, Sylvie Sanschagrin, Julie Champagne, David Juck, Kenneth Lee, and Charles W. Greer. 2015. “Microbial Community Composition, Functions, and Activities in the Gulf of Mexico 1 Year after the Deepwater Horizon Accident.” Edited by J. E. Kostka. *Applied and Environmental Microbiology* 81 (17): 5855–66. <https://doi.org/10.1128/AEM.01470-15>.
- Zaiko, Anastasija, Susanna A. Wood, Xavier Pochon, Laura Biessy, Olivier Laroche, Peter Croot, and Eva Garcia-Vazquez. 2020. “Elucidating Biodiversity Shifts in Ballast Water Tanks during a Cross-Latitudinal Transfer: Complementary Insights from Molecular Analyses.” *Environmental Science & Technology* 54 (13): 8443–54. <https://doi.org/10.1021/acs.est.0c01931>.
- Zhang, Shuangfei, Zhong Hu, and Hui Wang. 2018. “A Retrospective Review of Microbiological Methods Applied in Studies Following the Deepwater Horizon Oil Spill.” *Frontiers in Microbiology* 9 (March): 520. <https://doi.org/10.3389/fmicb.2018.00520>.
- Zhao, Beilun, Peter M. van Bodegom, and Krijn Trimbos. 2022. “Environmental DNA Methylation of *Lymnaea stagnalis* Varies with Age and Is Hypermethylated Compared to Tissue DNA.” *Molecular Ecology Resources* 23 (1). <https://doi.org/10.1111/1755-0998.13691>.
- Zimmermann, Jonas, Gernot Glöckner, Regine Jahn, Neela Enke, and Birgit Gemeinholzer. 2014. “Metabarcoding vs. Morphological Identification to Assess Diatom Diversity in Environmental Studies.” *Molecular Ecology Resources* 15 (3): 526–42. <https://doi.org/10.1111/1755-0998.12336>.
- Zizka, Vera M.A., Florian Leese, Bianca Peinert, and Matthias F. Geiger. 2019. “DNA Metabarcoding from Sample Fixative as a Quick and Voucher-Preserving Biodiversity Assessment Method.” *Genome* 62 (3): 122–36. <https://doi.org/10.1139/gen-2018-0048>.